

Product datasheet

Anti-ATP5H antibody [7F9BG1] ab110275

★★★★★ 3 Abreviews 11 References 4 Images

Overview

Product name	Anti-ATP5H antibody [7F9BG1]
Description	Mouse monoclonal [7F9BG1] to ATP5H
Host species	Mouse
Tested applications	Suitable for: WB, ICC/IF, Flow Cyt
Species reactivity	Reacts with: Mouse, Rat, Cow, Human, African green monkey
Immunogen	Other Immunogen Type corresponding to Cow ATP5H.
Positive control	Isolated mitochondria from Human heart, Bovine heart, Rat heart, Mouse heart, and HepG2, Cultured Human embryonic lung-derived fibroblasts (strain MRC5), HeLa cells
General notes	<p>This antibody clone is manufactured by Abcam.</p> <p>For mouse and rat samples, this antibody will only recognize ATP5H in purified mitochondrial samples. Mouse and rat cell lysates and tissue homogenates are not recommended with this antibody.</p> <p>Product was previously marketed under the MitoSciences sub-brand.</p> <p>If you require this antibody in a particular buffer formulation or a particular conjugate for your experiments, please contact orders@abcam.com or you can find further information here.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze.
Storage buffer	Preservative: 0.02% Sodium azide Constituent: HEPES buffered saline
Purity	Immunogen affinity purified
Purification notes	Near homogeneity as judged by SDS-PAGE. ab110275 was produced in vitro using hybridomas grown in serum-free medium, and then purified by biochemical fractionation.
Clonality	Monoclonal
Clone number	7F9BG1
Isotype	IgG2b
Light chain type	kappa

Applications

Our [Abpromise guarantee](#) covers the use of **ab110275** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★★	Use a concentration of 1 µg/ml. Predicted molecular weight: 18 kDa.
ICC/IF		Use a concentration of 1 - 5 µg/ml. (heat-induced antigen-retrieval improves signal)
Flow Cyt	★★★★★	Use a concentration of 1 µg/ml. ab170192 - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.

Target

Function

Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core, and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F(0) domain and the peripheral stalk, which acts as a stator to hold the catalytic alpha(3)beta(3) subcomplex and subunit a/ATP6 static relative to the rotary elements.


Sequence similarities

Belongs to the ATPase d subunit family.

Cellular localization

Mitochondrion. Mitochondrion inner membrane.

Images



All lanes : Anti-ATP5H antibody [7F9BG1] (ab110275) at 1 µg/ml

Lane 1 : HDFn (human) cell lysates at 20 µg

Lane 2 : COS7 (monkey) cell lysates at 20 µg

Lane 3 : H4IIE (rat) cell lysates at 20 µg


Lane 4 : MEF (mouse) cell lysates at 20 µg

Lane 5 : bovine heart mitochondria lysates at 5 µg

Predicted band size: 18 kDa

Western blot - Anti-ATP5H antibody [7F9BG1]

(ab110275)



Western blot - Anti-ATP5H antibody [7F9BG1] (ab110275)

All lanes : Anti-ATP5H antibody [7F9BG1] (ab110275) at 1/1 dilution

Lane 1 : Human heart mitochondria at 10 µg


Lane 2 : Bovine heart mitochondria at 4 µg

Lane 3 : Rat heart mitochondria at 10 µg

Lane 4 : Mouse heart mitochondria at 10 µg


Lane 5 : HepG2 mitochondria at 20 µg

Predicted band size: 18 kDa



Immunocytochemistry/ Immunofluorescence - Anti-ATP5H antibody [7F9BG1] (ab110275)

Mitochondrial localization of ATP5H. Cultured Human embryonic lung-derived fibroblasts (strain MRC5) were fixed, permeabilized and then labeled with ab110275 (5 µg/ml) followed by an AlexaFluor® 488-conjugated-goat-anti-mouse IgG(H+L) secondary antibody (2 µg/ml).



Flow Cytometry - Anti-ATP5H antibody [7F9BG1] (ab110275)

HeLa cells were stained with 1 µg/mL ab110275 (blue) or an equal amount of an isotype control antibody (red) and analyzed by flow cytometry.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors