

Product datasheet

Anti-CD22 antibody [MEM-01], prediluted (Phycoerythrin) ab27548

1 Image

Overview

Product name	Anti-CD22 antibody [MEM-01], prediluted (Phycoerythrin)
Description	Mouse monoclonal [MEM-01] to CD22, prediluted (Phycoerythrin)
Host species	Mouse
Conjugation	Phycoerythrin. Ex: 488nm, Em: 575nm
Specificity	ab27548 recognises CD22.
Tested applications	Suitable for: Flow Cyt
Species reactivity	Reacts with: Human Predicted to work with: Non human primates ▲
Immunogen	Burkitt's lymphoma cell line Raji.

Properties

Form	Prediluted
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.097% Sodium azide Constituent: 0.2% BSA High grade protease free BSA
Purity	IgG fraction
Clonality	Monoclonal
Clone number	MEM-01
Isotype	IgG1

Applications

Our [Abpromise guarantee](#) covers the use of **ab27548** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
-------------	-----------	-------

Flow Cyt

Application notes

Flow Cyt: Use 20µl for 10⁶ cells or 100µl of whole blood.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

Target

Function

Mediates B-cell B-cell interactions. May be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B-cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules.

Tissue specificity

B-lymphocytes.

Sequence similarities

Belongs to the immunoglobulin superfamily. SIGLEC (sialic acid binding Ig-like lectin) family.

Contains 6 Ig-like C2-type (immunoglobulin-like) domains.

Contains 1 Ig-like V-type (immunoglobulin-like) domain.

Domain

Contains 4 copies of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.

Post-translational modifications


Phosphorylation of Tyr-762, Tyr-807 and Tyr-822 are involved in binding to SYK, GRB2 and SYK, respectively. Phosphorylation of Tyr-842 is involved in binding to SYK, PLCG2 and PIK3R1/PIK3R2.

Phosphorylated on tyrosine residues by LYN.

Cellular localization

Cell membrane.

Images



ab27548 staining CD22 on human peripheral blood cells. Cell population was gated on lymphocytes.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors