

Product datasheet

Anti-CD86 antibody [GL-1] ab119857

1 Abreviews 8 References

Overview

Product name	Anti-CD86 antibody [GL-1]
Description	Rat monoclonal [GL-1] to CD86
Host species	Rat
Tested applications	Suitable for: IHC-Fr, Functional Studies, Flow Cyt, Blocking, ICC/IF, IP
Species reactivity	Reacts with: Mouse
Immunogen	LPS-activated CBA/Cs Mouse splenic B cells.
Positive control	Mouse splenocytes.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C. Do Not Freeze. Store undiluted.
Storage buffer	pH: 7.20 Preservative: 0.09% Sodium azide Constituents: 0.87% Sodium chloride, 0.14% Monobasic dihydrogen sodium phosphate
Purity	Affinity purified
Purification notes	This purified format is guaranteed to be >90% pure as determined by SDS-PAGE analysis.
Clonality	Monoclonal
Clone number	GL-1
Isotype	IgG2a

Applications

Our [Abpromise guarantee](#) covers the use of **ab119857** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-Fr		Use at an assay dependent concentration.

Application	Abreviews	Notes
Functional Studies		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. ab18450 - Rat monoclonal IgG2a, is suitable for use as an isotype control with this antibody.
Blocking		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.

Target

Function	Receptor involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. May play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. Isoform 2 interferes with the formation of CD86 clusters, and thus acts as a negative regulator of T-cell activation.
Tissue specificity	Expressed by activated B-lymphocytes and monocytes.
Sequence similarities	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
Post-translational modifications	Polyubiquitinated; which is promoted by MARCH8 and results in endocytosis and lysosomal degradation.
Cellular localization	Membrane.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors