

Product datasheet

Anti-DRAK2 antibody ab8419

1 Abreviews 3 Images

Overview

Product name	Anti-DRAK2 antibody
Description	Rabbit polyclonal to DRAK2
Host species	Rabbit
Tested applications	Suitable for: WB, ICC/IF
Species reactivity	Reacts with: Human
Immunogen	Synthetic peptide: CSKRFRFDDSLPNPHE , corresponding to amino acids 351/365 of Human DRAK2. (Peptide available as ab8454 .)

 [Run BLAST with](#)  [Run BLAST with](#)

Positive control Wholecell lysate from Jurkat cells An approximately 45 kDa band can be detected.

General notes

Apoptosis is mediated by death domain containing adapter molecules and a caspase family of proteases. Certain serine/threonine protein kinases, such as ASK-1 and RIP, are mediators of apoptosis. Two novel serine/threonine kinases that induce apoptosis were recently identified and designated DRAK1 and DRAK2 (for DAP kinase-related apoptosis-inducing protein kinases) (1). DRAKs contain an N-terminal kinase domain and a C-terminal regulation domain. Overexpression of DRAK2 induces apoptosis. DRAKs have high sequence homology to DAP and ZIP kinases, and they represent a novel family of serine/threonine kinases, which mediates apoptosis through their catalytic activities. DRAK2 is located in nucleus and the messenger RNA was ubiquitously expressed in human tissues.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	PBS with 0.02% sodium azide
Purity	Ion Exchange Chromatography
Purification notes	DRAK2 Antibody is Ion exchange chromatography purified.
Primary antibody notes	Apoptosis is mediated by death domain containing adapter molecules and a caspase family of proteases. Certain serine/threonine protein kinases, such as ASK-1 and RIP, are mediators of apoptosis. Two novel serine/threonine kinases that induce apoptosis were recently identified

and designated DRAK1 and DRAK2 (for DAP kinase-related apoptosis-inducing protein kinases) (1). DRAKs contain an N-terminal kinase domain and a C-terminal regulation domain. Overexpression of DRAK2 induces apoptosis. DRAKs have high sequence homology to DAP and ZIP kinases, and they represent a novel family of serine/threonine kinases, which mediates apoptosis through their catalytic activities. DRAK2 is located in nucleus and the messenger RNA was ubiquitously expressed in human tissues.

Clonality	Polyclonal
Isotype	IgG
Light chain type	unknown

Applications

Our [Abpromise guarantee](#) covers the use of **ab8419** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/1000. Detects a band of approximately 45 kDa (predicted molecular weight: 43 kDa). Can be blocked with Human DRAK2 peptide (ab8454) .
ICC/IF		Use a concentration of 10 µg/ml.

Target

Function	Acts as a positive regulator of apoptosis.
Tissue specificity	Highly expressed in placenta, lung, pancreas. Lower levels in heart, brain, liver, skeletal muscle and kidney.
Sequence similarities	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. DAP kinase subfamily. Contains 1 protein kinase domain.
Post-translational modifications	Autophosphorylated.
Cellular localization	Nucleus.

Images




Western blot - Anti-DRAK2 antibody (ab8419)

Lane 1 : Anti-DRAK2 antibody (ab8419) at 1 $\mu\text{g/ml}$ (DRAK2 antibody)

Lane 2 : Anti-DRAK2 antibody (ab8419) at 2 $\mu\text{g/ml}$ (DRAK2 antibody)

All lanes : Raji cell lysate

Predicted band size: 43 kDa



Western blot - Anti-DRAK2 antibody (ab8419)

All lanes : Anti-DRAK2 antibody (ab8419) at 1/500 dilution

Lane 1 : Jurkat whole cell lysate with absence of blocking peptide

Lane 2 : Raji whole cell lysate with absence of blocking peptide


Lane 3 : Jurkat whole cell lysate with presence of blocking peptide

Lane 4 : Raji whole cell lysate with presence of blocking peptide

Predicted band size: 43 kDa

Observed band size: 45 kDa

We are unsure about the nature of the 70 kDa band. However, DRAK2 is autophosphorylated and it is possible that this band corresponds to the phosphorylated form. The fact that the detection of this band is blocked by DRAK2 peptide indicates that it probably is closely related to the DRAK2 protein.



ab8419 at 10µg/ml staining DRAK2 in Jurkat cells by ICC/IF

Immunocytochemistry/ Immunofluorescence - Anti-DRAK2 antibody (ab8419)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors