


## Product datasheet

# Anti-Frizzled 2 antibody ab52565

### 3 References

#### Overview

<b>Product name</b>	Anti-Frizzled 2 antibody
<b>Description</b>	Rabbit polyclonal to Frizzled 2
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> WB, IP
<b>Species reactivity</b>	<b>Reacts with:</b> Human <b>Predicted to work with:</b> Mouse 
<b>Immunogen</b>	Synthetic peptide derived from the N-terminal region of human FZD2.
<b>Positive control</b>	Human PC-3, SKOV3, SW480 and HEK293 cell lysates for Western blotting and HepG2 for immunoprecipitation.

#### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
<b>Storage buffer</b>	Preservative: 0.1% Sodium Azide Constituents: PBS, pH 7.4
<b>Purity</b>	Immunogen affinity purified
<b>Purification notes</b>	Purified from Rabbit antiserum
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG

#### Applications

Our [Abpromise guarantee](#) covers the use of **ab52565** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes

Application	Abreviews	Notes
WB		Use a concentration of 1 - 3 µg/ml. Detects a band of approximately 85 kDa (predicted molecular weight: 63 kDa). The observed molecular weight of ~ 85 kDa may be due to post-translational modifications such as glycosylation, which is predicted in the amino acid sequence.
IP		Use at an assay dependent dilution. Use at a concentration of 7 µg/reaction.

## Target

<b>Function</b>	Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.
<b>Tissue specificity</b>	Widely expressed. In the adult, mainly found in heart, placenta, skeletal muscle, lung, kidney, pancreas, prostate, testis, ovary and colon. In the fetus, expressed in brain, lung and kidney. Low levels in fetal liver.
<b>Sequence similarities</b>	Belongs to the G-protein coupled receptor Fz/Smo family. Contains 1 FZ (frizzled) domain.
<b>Domain</b>	Lys-Thr-X-X-Trp motif is involved in the activation of the Wnt/beta-catenin signaling pathway. The FZ domain is involved in binding with Wnt ligands.
<b>Cellular localization</b>	Membrane.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors