

Product datasheet

Anti-GATA4 antibody [EPR4768] ab134057

Recombinant RabMAB

★★★★☆ 2 Abreviews 5 References 8 Images

Overview

Product name	Anti-GATA4 antibody [EPR4768]
Description	Rabbit monoclonal [EPR4768] to GATA4
Host species	Rabbit
Tested applications	Suitable for: WB, Flow Cyt, ICC/IF Unsuitable for: IHC-P or IP
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide corresponding to Human GATA4 aa 1-100 (N terminal). Database link: P43694
Positive control	HepG2, Caco-2 and HeLa cell lysates; HepG2 cells.
General notes	This antibody was developed as part of a collaboration between Dartmouth College and the lab of Sergei Tevosian. A trial size is available to purchase for this antibody. Our RabMAB [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMab[®] patents We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team. This product is a recombinant rabbit monoclonal antibody.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
Dissociation constant (K_D)	K _D = 2.00 x 10 ⁻¹² M





[Learn more about K_D](#)

Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
Purity	Immunogen affinity purified
Clonality	Monoclonal
Clone number	EPR4768
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab134057** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★☆	1/1000 - 1/10000. Predicted molecular weight: 54 kDa.

Flow Cyt 1/600.
For unpurified use at 1/1200.

[ab172730](#) - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

ICC/IF ★★★★★
For unpurified use at 1/100 - 1/250.

Application notes Is unsuitable for IHC-P or IP.

Target

Function Transcriptional activator that binds to the consensus sequence 5'-AGATAG-3' and plays a key role in cardiac development (PubMed:24000169). Involved in bone morphogenetic protein (BMP)-mediated induction of cardiac-specific gene expression (By similarity). Binds to BMP response element (BMPRE) DNA sequences within cardiac activating regions (By similarity). Acts as a transcriptional activator of ANF in cooperation with NKX2-5 (By similarity). Promotes cardiac myocyte enlargement (PubMed:20081228). Required during testicular development (PubMed:21220346). May play a role in sphingolipid signaling by regulating the expression of sphingosine-1-phosphate degrading enzyme, sphingosine-1-phosphate lyase (PubMed:15734735).

Involvement in disease Atrial septal defect 2
Ventricular septal defect 1

Tetralogy of Fallot

Atrioventricular septal defect 4

Testicular anomalies with or without congenital heart disease

GATA4 mutations can predispose to dilated cardiomyopathy (CMD), a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death.

Sequence similarities

Contains 2 GATA-type zinc fingers.


Post-translational modifications

Methylation at Lys-300 attenuates transcriptional activity.


Cellular localization

Nucleus.


Images



Overlay histogram showing Caco-2 cells stained with ab134057 (red line) at 1/600 dilution. The cells were fixed with 2% paraformaldehyde. The secondary antibody used was a FITC conjugated goat anti-rabbit IgG at 1/150 dilution. Isotype control antibody (green line) was rabbit monoclonal IgG used under the same conditions.



ab134057 staining GATA4 in the PC-3 cell line by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100. Samples were incubated with primary antibody (1/150). An Alexa Fluor[®]555-conjugated Goat anti-rabbit IgG(1/500) was used as the secondary antibody. Nuclei were counterstained with DAPI.




Western blot - Anti-GATA4 antibody [EPR4768]
(ab134057)

Anti-GATA4 antibody [EPR4768] (ab134057)
at 1/2000 dilution + Mouse spleen lysate at 10
 μ g

Secondary

Goat Anti-Rabbit IgG, (H+L), HRP-conjugated
at 1/1000 dilution

Predicted band size: 54 kDa




Western blot - Anti-GATA4 antibody [EPR4768]
(ab134057)

Anti-GATA4 antibody [EPR4768] (ab134057)
at 1/2000 dilution + HepG2 lysate at 10 μ g

Secondary


Goat Anti-Rabbit IgG, (H+L), HRP-conjugated

Predicted band size: 54 kDa



Flow Cytometry - Anti-GATA4 antibody [EPR4768] (ab134057)

Overlay histogram showing HeLa cells stained with ab134057, unpurified (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab134057, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HeLa cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Western blot - Anti-GATA4 antibody [EPR4768] (ab134057)

All lanes : Anti-GATA4 antibody [EPR4768] (ab134057) at 1/1000 dilution (unpurified)

Lane 1 : HepG2 cell lysate

Lane 2 : Caco-2 cell lysate


Lane 3 : HeLa cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 54 kDa



Immunofluorescent analysis of HepG2 cells labelling GATA4 with ab134057, unpurified, at 1/100 dilution.

Immunocytochemistry/ Immunofluorescence - Anti-GATA4 antibody [EPR4768] (ab134057)



Equilibrium disassociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Other - Anti-GATA4 antibody [EPR4768] (ab134057)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors