

Product datasheet

Human TEM8 peptide ab41852

Overview

Product name Human TEM8 peptide

Description

Nature Synthetic

Amino Acid Sequence

Species Human

Specifications

Our [Abpromise guarantee](#) covers the use of **ab41852** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications Blocking - Blocking peptide for Anti-TEM8 antibody ([ab21270](#))

Form Liquid

Preparation and Storage

Stability and Storage Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Preservative: 0.02% Sodium Azide

Constituents: 0.1% BSA, PBS, pH 7.2

General Info

Function Plays a role in cell attachment and migration. Interacts with extracellular matrix proteins and with the actin cytoskeleton. Mediates adhesion of cells to type 1 collagen and gelatin, reorganization of the actin cytoskeleton and promotes cell spreading. Plays a role in the angiogenic response of cultured umbilical vein endothelial cells.

Tissue specificity Detected in umbilical vein endothelial cells (at protein level). Highly expressed in tumor endothelial cells.

Involvement in disease Defects in ANTXR1 are associated with susceptibility to hemangioma capillary infantile (HCI) [MIM:602089]. HCI are benign, highly proliferative lesions involving aberrant localized growth of capillary endothelium. They are the most common tumor of infancy, occurring in up to 10% of all births. Hemangiomas tend to appear shortly after birth and show rapid neonatal growth for up to

12 months characterized by endothelial hypercellularity and increased numbers of mast cells. This phase is followed by slow involution at a rate of about 10% per year and replacement by fibrofatty stroma.

Sequence similarities

Belongs to the ATR family.
Contains 1 VWFA domain.

Domain

Binding to PA occurs through the VWA domain.

Cellular localization

Cell membrane. Cell projection > lamellipodium membrane. Cell projection > filopodium membrane. At the membrane of lamellipodia and at the tip of actin-enriched filopodia.
Colocalizes with actin at the base of lamellipodia.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors