

## Product datasheet

# Anti-MICB antibody [MM0473-3C37] ab90237

### Overview

<b>Product name</b>	Anti-MICB antibody [MM0473-3C37]
<b>Description</b>	Mouse monoclonal [MM0473-3C37] to MICB
<b>Host species</b>	Mouse
<b>Tested applications</b>	<b>Suitable for:</b> WB, Flow Cyt
<b>Species reactivity</b>	<b>Reacts with:</b> Human
<b>Immunogen</b>	Recombinant full length Human MICB.

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
<b>Storage buffer</b>	Constituent: PBS
<b>Purity</b>	Protein G purified
<b>Purification notes</b>	The IgG fraction of culture supernatant was purified by Protein G affinity chromatography and 0.2 µm filtered.
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	MM0473-3C37
<b>Isotype</b>	IgG2b

### Applications

Our [Abpromise guarantee](#) covers the use of **ab90237** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/500 - 1/1000. Predicted molecular weight: 43 kDa.
Flow Cyt		1/20 - 1/200. <a href="#">ab170192</a> - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.

## Target

---

<b>Function</b>	Seems to have no role in antigen presentation. Acts as a stress-induced self-antigen that is recognized by gamma delta T cells. Ligand for the KLRK1/NKG2D receptor. Binding to KLRK1 leads to cell lysis.
<b>Tissue specificity</b>	Widely expressed with the exception of the central nervous system where it is absent. Expressed in many, but not all, epithelial tumors of lung, breast, kidney, ovary, prostate and colon. In hepatocellular carcinomas, expressed in tumor cells but not in surrounding non-cancerous tissue.
<b>Involvement in disease</b>	Genetic variations in MICA are a cause of susceptibility to rheumatoid arthritis (RA) [MIM:180300]. It is a systemic inflammatory disease with autoimmune features and a complex genetic component. It primarily affects the joints and is characterized by inflammatory changes in the synovial membranes and articular structures, widespread fibrinoid degeneration of the collagen fibers in mesenchymal tissues, and by atrophy and rarefaction of bony structures. Note=The MICB*004 allele is associated with rheumatoid arthritis. Note=Genetic variation in MICB is associated with cytomegalovirus and herpes simplex virus I seropositivity and this may be associated with schizophrenia risk.
<b>Sequence similarities</b>	Belongs to the MHC class I family. MIC subfamily. Contains 1 Ig-like C1-type (immunoglobulin-like) domain.
<b>Post-translational modifications</b>	Proteolytically cleaved and released from the cell surface of tumor cells.
<b>Cellular localization</b>	Cell membrane. Binding to human cytomegalovirus glycoprotein UL16 causes sequestration in the endoplasmic reticulum.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors