

Product datasheet

Recombinant Human COMP protein ab124873

Overview

Product name	Recombinant Human COMP protein
Protein length	Full length protein

Description

Nature	Recombinant
Source	HEK 293 cells

Amino Acid Sequence

Accession [P49747](#)

Species Human

Sequence **HVDYKDDDDK**PAGQGQSPLGSDLGPQMLRELQETNAALQDVRELLRQQVR
 EITFLKNTVMECDACGMQQSVRTGLPSVRPQGQSPLGSDLGPQMLRELQE
 TNAALQDVRELLRQQVREITFLKNTVMECDACGMQQSVRTGLPSVRPLH
 CAPGFCFPGVACIQTESGARCGPCPAGFTGNGSHCTDVNECNAHPCFPRV
 RCINTSPGFRCEACPPGYSGPTHQGVGLAFKANKQVCTDINECETGQHN
 CVPNSVCINTRGSFQCGPCQPGFVGDQASGCQRRARFCPDGSPSECEH
 ADCVLERDGSRSVCVAVGWAGNGLCGRDIDLDFPDEKLRCPERQCRKD
 NCVTVPNSGQEDVDRDGIGDACPDPADGDGVPNEKDNCPLVRNPQQRNTD
 EDKWGDACDNCRSQKNDDQKDTDQDGRGDACDDDDIDGDRIRNQADNCPRV
 PNSDQKDSGDGIGDACDNCQKSNPDQADVDHDFVGDACDSDQDQDGDG
 HQDSRDNCPTVPNSAQEDSDHDGQGDACDDDDNDGVPDSRDNCRLVNP
 GQEDADRDRGVGDVCQDDFDADKVVDKIDVCPENAEVTLDFRAFQTVVLD
 PEGDAQIDPNWVVLNQGREGVQTMNSDPGLAVGYTAFNGVDFEGTFHVNT
 VTDDDYAGFIFGYQDSSSFYVVMWKQMEQTYWQANPFRAVAEPGIQLKAV
 KSSTGPGEQLRNALWHTGDTESQVRLLWKDPRNVGWKDKKSYRWFLQHRP
 QVGYIRVRFYEGPELVADSNVLDTTMRGGRLGVFCFSQENIIVANLRYR
 CNDTIPEDYETHQLRQA

Molecular weight	82 kDa including tags
Amino acids	21 to 757
Tags	DDDDK tag N-Terminus

Specifications

Our [Abpromise guarantee](#) covers the use of **ab124873** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE Western blot
Form	Lyophilised

Preparation and Storage

Stability and Storage	Shipped at 4°C. Store at -20°C. pH: 7.50 Constituents: 0.24% Tris, 0.29% Sodium chloride
------------------------------	--

Reconstitution	It is recommended to add deionized water to prepare a working stock solution of approximately 0.5mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it on cell culture. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles.
-----------------------	---

General Info

Function	May play a role in the structural integrity of cartilage via its interaction with other extracellular matrix proteins such as the collagens and fibronectin. Can mediate the interaction of chondrocytes with the cartilage extracellular matrix through interaction with cell surface integrin receptors. Could play a role in the pathogenesis of osteoarthritis. Potent suppressor of apoptosis in both primary chondrocytes and transformed cells. Suppresses apoptosis by blocking the activation of caspase-3 and by inducing the IAP family of survival proteins (BIRC3, BIRC2, BIRC5 and XIAP). Essential for maintaining a vascular smooth muscle cells (VSMCs) contractile/differentiated phenotype under physiological and pathological stimuli. Maintains this phenotype of VSMCs by interacting with ITGA7.
Tissue specificity	Abundantly expressed in the chondrocyte extracellular matrix, and is also found in bone, tendon, ligament and synovium and blood vessels. Increased amounts are produced during late stages of osteoarthritis in the area adjacent to the main defect.
Involvement in disease	Defects in COMP are the cause of multiple epiphyseal dysplasia type 1 (EDM1) [MIM:132400]. EDM is a generalized skeletal dysplasia associated with significant morbidity. Joint pain, joint deformity, waddling gait, and short stature are the main clinical signs and symptoms. EDM is broadly categorized into the more severe Fairbank and the milder Ribbing types. Defects in COMP are the cause of pseudoachondroplasia (PSACH) [MIM:177170]. PSACH is a dominantly inherited chondrodysplasia characterized by short stature and early-onset osteoarthritis. PSACH is more severe than EDM1 and is recognized in early childhood.
Sequence similarities	Belongs to the thrombospondin family. Contains 4 EGF-like domains. Contains 1 TSP C-terminal (TSPC) domain. Contains 8 TSP type-3 repeats.
Developmental stage	Present during the earliest stages of limb maturation and is later found in regions where the joints develop.
Domain	The cell attachment motif mediates the attachment to chondrocytes. It mediates the induction of both the IAP family of survival proteins and the antiapoptotic response. The TSP C-terminal domain mediates interaction with FN1 and ACAN.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors