

## Product datasheet

# Recombinant Human Histamine H1 Receptor co-expressed with RGS4 protein ab90479

### Overview

---

<b>Product name</b>	Recombinant Human Histamine H1 Receptor co-expressed with RGS4 protein
<b>Protein length</b>	Full length protein

### Description

---

<b>Nature</b>	Recombinant
<b>Source</b>	Baculovirus infected Sf9 cells
<b>Amino Acid Sequence</b>	
<b>Species</b>	Human

### Specifications

---

Our [Abpromise guarantee](#) covers the use of **ab90479** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>Applications</b>	Functional Studies
<b>Form</b>	Liquid

### Preparation and Storage

---

<b>Stability and Storage</b>	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 7.40 Constituents: 0.11875% Magnesium chloride, 1.185% Tris HCl, 0.0292% EDTA
------------------------------	---

### General Info

---

<b>Relevance</b>	Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like cells, and neurons. The histamine H1 receptor is an integral membrane protein and belongs to the family 1 of G protein-coupled receptors. It mediates the contraction of smooth muscles, increase in capillary permeability due to contraction of terminal venules, catecholamine release from adrenal medulla, and mediates neurotransmission in the central nervous system. Regulator of G protein signaling 4 (RGS4) negatively regulates G protein coupled receptor signaling. RGS4 belongs to a family of regulatory molecules that act as GTPase activating proteins (GAPs) for G
------------------	--

alpha subunits of heterotrimeric G proteins. RGS proteins are able to deactivate G protein subunits of the Gi alpha, Go alpha and Gq alpha subtypes. They drive G proteins into their inactive GDP bound forms. All RGS proteins share a conserved 120 amino acid sequence termed the RGS domain. Regulator of G protein signaling 4 protein is 37% identical to RGS1 and 97% identical to rat Rgs4. This protein negatively regulate signaling upstream or at the level of the heterotrimeric G protein and is localized in the cytoplasm. The antihypertrophic effects of RGS4 in the myocardium is well characterized. RGS4 may also have a potential role in the pathogenesis of cardiac arrhythmias.

#### **Cellular localization**

Histamine H1 Receptor: Cell membrane; Multi-pass membrane protein.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

#### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors