

Product datasheet

Recombinant Human ING2 protein (denatured) ab171499

[1 Image](#)

Overview

Product name	Recombinant Human ING2 protein (denatured)
Protein length	Full length protein
Description	Recombinant Human ING2 protein

Description

Nature	Recombinant
Source	Escherichia coli
Amino Acid Sequence	
Accession	Q9H160
Species	Human

Sequence	<p>MGSSHHHHHH SSGLVPRGSH MGSMLGQQQQ QLYSSAALLT GERSRLLTCYVQDYLECVES LPHDMQRNVS VLRELDNKYQ ETLKEIDVY EKYKKEDDLN QKKRLQQLLQ RALINSQELG DEKIQV/TQM LELVENRARQ MELHSQCFQD PAESERASDK AKMDSSQPER SSRRPRRQRT SESRDLCHMA NGIEDCDDQP PKEKKSksak KKKRSKAKQE REASPVEFAIDPNEPTYCLC NQVSYGEMIG CDNEQCPIEW FHFSCVSLTY KPKGKWCYCPK CRGDNEKTMD KSTKTKKDR RSR</p>
-----------------	--

Molecular weight	35 kDa including tags
Amino acids	1 to 280
Tags	His tag N-Terminus

Specifications

Our [Abpromise guarantee](#) covers the use of **ab171499** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications	SDS-PAGE
Purity	> 85 % SDS-PAGE.

ab171499 is purified by using anion-exchange chromatography (DEAE sepharose resin) and gel-filtration chromatography.

Form Liquid

Preparation and Storage

Stability and Storage Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

pH: 8.00

Constituents: 2.4% Urea, 0.32% Tris HCl, 10% Glycerol

General Info

Function Seems to be involved in p53/TP53 activation and p53/TP53-dependent apoptotic pathways, probably by enhancing acetylation of p53/TP53. Component of a mSin3A-like corepressor complex, which is probably involved in deacetylation of nucleosomal histones. ING2 activity seems to be modulated by binding to phosphoinositides (PtdInsPs).


Tissue specificity Widely expressed. Higher expressed in colon-cancer tumor than in normal colon tissues.

Sequence similarities Belongs to the ING family.
Contains 1 PHD-type zinc finger.

Domain The PHD-type zinc finger domain binds to phosphoinositides (PtdInsPs), including phosphatidylinositol 5-phosphate (PtdIns(5)P).

Cellular localization Nucleus. Predominantly nuclear. Localized to chromatin and nuclear matrix. Upon reduced PtdIns(5)P levels seems to be released from chromatin and, at least partially, translocated to the cytoplasm.

Images



15% SDS-PAGE analysis of ab171499 (3µg).

SDS-PAGE - Recombinant Human ING2 protein
(denatured) (ab171499)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors