

Product datasheet

Recombinant Human Lipocalin-2 / NGAL protein ab92916

[1 Image](#)

Overview

Product name Recombinant Human Lipocalin-2 / NGAL protein
Protein length Full length protein

Description

Nature Recombinant
Source Escherichia coli
Amino Acid Sequence
Species Human
Amino acids 19 to 198

Specifications

Our [Abpromise guarantee](#) covers the use of **ab92916** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Applications Western blot
 ELISA
 Mass Spectrometry
 SDS-PAGE
Form Lyophilised
Additional notes After reconstitution, store in working aliquots at -80°C.

Preparation and Storage

Stability and Storage Shipped at 4°C. Store at 4°C prior to reconstitution. Upon delivery aliquot. Store at -80°C.
 Reconstitute for long term storage.
 Preservative: None
 Constituents: PBS, pH 7.4

General Info

Function

Iron-trafficking protein involved in multiple processes such as apoptosis, innate immunity and renal development. Binds iron through association with 2,5-dihydroxybenzoic acid (2,5-DHBA), a siderophore that shares structural similarities with bacterial enterobactin, and delivers or removes iron from the cell, depending on the context. Iron-bound form (holo-24p3) is internalized following binding to the SLC22A17 (24p3R) receptor, leading to release of iron and subsequent increase of intracellular iron concentration. In contrast, association of the iron-free form (apo-24p3) with the SLC22A17 (24p3R) receptor is followed by association with an intracellular siderophore, iron chelation and iron transfer to the extracellular medium, thereby reducing intracellular iron concentration. Involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular iron concentration without promoting apoptosis, while iron-free form decreases intracellular iron levels, inducing expression of the proapoptotic protein BCL2L11/BIM, resulting in apoptosis. Involved in innate immunity, possibly by sequestering iron, leading to limit bacterial growth.

Tissue specificity

Expressed in bone marrow and in tissues that are prone to exposure to microorganism. High expression is found in bone marrow as well as in uterus, prostate, salivary gland, stomach, appendix, colon, trachea and lung. Not found in the small intestine or peripheral blood leukocytes.


Sequence similarities

Belongs to the calycin superfamily. Lipocalin family.

Cellular localization

Secreted. Upon binding to the SLC22A17 (24p3R) receptor, it is internalized.

Images



Western blot - Recombinant Human Lipocalin-2 /
NGAL protein (ab92916)

ab92916 analysed on 10% SDS-PAGE and stained with
Coomassie Blue (1) or detected by western blot analysis with an
antibody to the proprietary tag (2).

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors