

Product datasheet

Anti-CD4 antibody [EDU-2] (Phycoerythrin) ab1155

Overview

Product name	Anti-CD4 antibody [EDU-2] (Phycoerythrin)
Description	Mouse monoclonal [EDU-2] to CD4 (Phycoerythrin)
Host species	Mouse
Conjugation	Phycoerythrin. Ex: 488nm, Em: 575nm
Tested applications	Suitable for: Flow Cyt
Species reactivity	Reacts with: Human
Immunogen	Human T-lymphocytes.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.1% Sodium azide Constituent: 0.5% BSA
Clonality	Monoclonal
Clone number	EDU-2
Myeloma	unknown
Isotype	IgG2a
Light chain type	unknown

Applications

Our [Abpromise guarantee](#) covers the use of **ab1155** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
-------------	-----------	-------

Flow Cyt

Application notes

1. Counting and identification of the T-helper/inducer lymphocyte subset in human lysed whole peripheral blood or mononuclear cells separated by density gradient.
2. Studies of HIV infection. Patients infected with HIV lose CD4+ and contemporarily increase

CD8+ lymphocyte subset

3. Characterization of subtypes of T-cell leukaemias and lymphomas.

CD4 (PE) immunofluorescence analysis can be performed on a flow cytometer equipped with an excitation source of 488nm and fitted with logarithmic amplifiers.

10µl of CD4 (PE) is sufficient for labelling of 1×10^6 cells.

Molecular F/P ratio: 1.0

Target

Function	Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell activation. Induces the aggregation of lipid rafts.
Sequence similarities	Contains 3 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
Post-translational modifications	Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.
Cellular localization	Cell membrane. Localizes to lipid rafts. Removed from plasma membrane by HIV-1 Nef protein that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation. Cell surface expression is also down-modulated by HIV-1 Envelope polyprotein gp160 that interacts with, and sequesters CD4 in the endoplasmic reticulum.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors