

## Product datasheet

# Anti-TCR gamma + TCR delta antibody [TCR1] (Biotin) ab25151

### Overview

---

<b>Product name</b>	Anti-TCR gamma + TCR delta antibody [TCR1] (Biotin)
<b>Description</b>	Mouse monoclonal [TCR1] to TCR gamma + TCR delta (Biotin)
<b>Host species</b>	Mouse
<b>Conjugation</b>	Biotin
<b>Tested applications</b>	<b>Suitable for:</b> IHC-Fr, Flow Cyt, IP
<b>Species reactivity</b>	<b>Reacts with:</b> Chicken
<b>Immunogen</b>	The details of the immunogen for this antibody are not available.
<b>General notes</b>	ab25151 can be used for in ovo depletion of gamma delta T cells.

### Properties

---

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	Preservative: 0.1% Sodium azide Constituent: PBS
<b>Purity</b>	IgG fraction
<b>Primary antibody notes</b>	ab25151 can be used for in ovo depletion of gamma delta T cells.
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	TCR1
<b>Isotype</b>	IgG1
<b>Light chain type</b>	kappa

### Applications

---

Our [Abpromise guarantee](#) covers the use of **ab25151** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-Fr		Use at an assay dependent concentration. Use with acetone fixed tissues.
Flow Cyt		Use 1µg for 10 <sup>6</sup> cells. <a href="#">ab18434</a> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
IP		Use at an assay dependent concentration.

## Target

### Relevance

T cell receptors (TCR) recognize foreign antigens which have been processed as small peptides and bound to major histocompatibility complex (MHC) molecules at the surface of antigen presenting cells (APC). Each T cell receptor is a dimer consisting of one α and one β chain or one δ and one γ chain. This region represents the germline organization of the T cell receptor beta locus. The beta locus includes V (variable), J (joining), diversity (D), and C (constant) segments. During T cell development, the beta chain is synthesized by a recombination event at the DNA level joining a D segment with a J segment; a V segment is then joined to the D-J gene. The C segment is later joined by splicing at the RNA level. The g/d TCR associates with CD3 and is expressed on a T cell subset found in the thymus, the intestinal epithelium, and the peripheral lymphoid tissues and peritoneum. Most g/d T cells are CD4-/CD8-, some are CD8+. T cells expressing the g/d TCR have been shown to play a role in oral tolerance, tumor-associated tolerance, and autoimmune disease.

### Cellular localization

Type I membrane protein

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors