

Product datasheet

Anti-CD4 antibody [74-12-4] (Phycoerythrin) ab25354

[2 Images](#)

Overview

Product name	Anti-CD4 antibody [74-12-4] (Phycoerythrin)
Description	Mouse monoclonal [74-12-4] to CD4 (Phycoerythrin)
Host species	Mouse
Conjugation	Phycoerythrin. Ex: 488nm, Em: 575nm
Specificity	ab25354 recognises porcine CD4a.
Tested applications	Suitable for: Flow Cyt, IP, Microcytotoxicity testing
Species reactivity	Reacts with: Pig
Immunogen	The details of the immunogen for this antibody are not available.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.09% Sodium Azide Constituents: 16% Sucrose, PBS; Stabilizing agent
Purity	IgG fraction
Clonality	Monoclonal
Clone number	74-12-4
Isotype	IgG2b
Light chain type	kappa

Applications

Our [Abpromise guarantee](#) covers the use of **ab25354** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		

Application	Abreviews	Notes
-------------	-----------	-------

IP

Microcytotoxicity testing

Application notes

Flow Cyt: Use 0.2µg for 10⁶ cells.

IP: Use at an assay dependent dilution.

Mct: Use at an assay dependent dilution.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

Target

Function

Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell activation. Induces the aggregation of lipid rafts.

Sequence similarities

Contains 3 Ig-like C2-type (immunoglobulin-like) domains.

Contains 1 Ig-like V-type (immunoglobulin-like) domain.


Post-translational modifications

Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.

Cellular localization


Cell membrane. Localizes to lipid rafts. Removed from plasma membrane by HIV-1 Nef protein that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation. Cell surface expression is also down-modulated by HIV-1 Envelope glycoprotein gp160 that interacts with, and sequesters CD4 in the endoplasmic reticulum.

Images



Flow Cytometry analysis of Porcine peripheral blood lymphocytes labeling CD4 with ab25354 at 0.1 µg/10⁶ cells. A Mouse Anti-Porcine CD8α-FITC was used as the secondary antibody.

Flow Cytometry - Anti-CD4 antibody [74-12-4]
(Phycoerythrin) (ab25354)



Flow Cytometry - Anti-CD4 antibody [74-12-4]
(Phycoerythrin) (ab25354)

Peripheral blood mononuclear cells were isolated from heparinized pig blood on Ficoll-Hypaque density gradients and double-stained with ab25354 at 0.2ug/10⁶ cells and mouse anti-pig CD8a-FITC. Lymphocytes were gated and analyzed on a FACScan® flow cytometer (BDIS, San Jose, CA).

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors