

Product datasheet

Human stomach tissue lysate (signet-ring cell carcinoma) - soluble fraction (male, 43 years) ab44447

Overview

<b>Product name</b>	Human stomach tissue lysate (signet-ring cell carcinoma) - soluble fraction (male, 43 years)
<b>General notes</b>	<p>Location: Lesser curvature. Diagnosis: Signet-ring cell carcinoma. Stage: II; T<sub>3</sub>N<sub>0</sub>M<sub>0</sub> Grade: n/a</p> <p>Gross findings: Ulcerative lesion located at the lesser curvature, ill demarcated. Tumor measures 7 cm in diameter. Cut section is firm and white. Paragastric lymph nodes: 6.</p> <p>Microscopic findings: Tissue sections show proliferation of ring cells. Mitotic figures are evident. The neoplastic cells have coarse, irregular nuclei and scanty cytoplasm. The tumor cells form groups and glands and invade the muscularis propria, subserosa, and serosa. Lymph nodes show 0 of 7 positive.</p> <p>This product belongs to a range of individual sets of lysates that were extracted from very specific tissues and are gender-, cancer subtype-, tumor grade- and cellular morphology-specific. They allow for consistency, reproducibility, and comparative studies from experiment to experiment, due to the existence of specific (matched) lysates that can be used as controls. Two extractions from each tissue were performed, extraction 1 yielding the soluble protein fraction and extraction 2 the insoluble protein fraction. Please find below a range of products that relate to ab44447: Extraction 1: soluble protein fraction <a href="#">ab44447</a> Human Stomach Tumor Tissue lysate <a href="#">ab44446</a> Human Stomach Normal Tissue lysate (matched) Extraction 2: insoluble protein fraction <a href="#">ab44449</a> Human Stomach Tumor Tissue lysate <a href="#">ab44448</a> Human Stomach Normal Tissue lysate (matched).</p>

Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	<p>pH: 7.40</p> <p>Constituents: PBS, 0.0001% Pepstatin A, 0.018% Sodium orthovanadate, 0.004% Sodium fluoride, 0.0001% Aprotinin, 0.0001% Leupeptin, 0.25% Sodium deoxycholate, 0.017% PMSF, 0.1% SDS, 0.029% EDTA</p>
<b>Purification notes</b>	Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors