

# Anti-BDNF antibody ab6201

★★★★★ [1 Abreviews](#) [47 References](#) [5 Images](#)

### Overview

---

<b>Product name</b>	Anti-BDNF antibody
<b>Description</b>	Rabbit polyclonal to BDNF
<b>Host species</b>	Rabbit
<b>Specificity</b>	By dot blots, less than 0.1% cross-reactivity against mouse NGF, recombinant human NT3 or NT4. Regarding use with brain lysates in WB: BDNF is very weakly expressed in whole brain (orders of magnitude lower), and extremely difficult to detect in BRAIN lysates. Thus, it cannot be compared to structural proteins or more common targets. It is recommended to use a positive control or spiked lysate control to optimize WB conditions.
<b>Tested applications</b>	<b>Suitable for:</b> Dot blot, ICC/IF, ELISA, Neutralising, WB, IHC-FoFr
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Horse, Human, Pig
<b>Immunogen</b>	Recombinant full length protein (Human).
<b>Positive control</b>	IHC-Fr: rat dorsal root ganglion (using a ganglion from an animal perfused with 2% (2% not the usual 4%) formaldehyde, with the addition of 15% picric acid, will reveal BDNF in many small diameter neurons. WB: dorsal root ganglia is recommended as the best positive control, but if not available, then both brain or spinal cord extracts can be used.
<b>General notes</b>	<p>This antibody has proven particularly useful for western blot, immunohistochemistry and biological inhibition. For immunohistochemistry, cell bodies are clearly stained as well as some nerve terminals in the dorsal horn of the rat spinal cord. The antiserum does not stain the finest nerve terminals.</p> <p>For BDNF, multiple WB bands are possible and expected. The human protein has 5 isoforms (precursors: 28 – 37 kDa) and can be glycosylated (Uniprot: <a href="http://www.uniprot.org/uniprot/P23560">http://www.uniprot.org/uniprot/P23560</a>). The mature form is expected at ~14 kDa (monomer) and the dimer at ~28 kDa.</p>

### Properties

---

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Add glycerol to a final volume of 50% for extra stability and aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	Constituent: 0.0268% PBS
<b>Purity</b>	Protein G purified
<b>Clonality</b>	Polyclonal

<b>Isotype</b>	IgG
<b>Light chain type</b>	unknown

## Applications


**The Abpromise guarantee** Our **Abpromise guarantee** covers the use of ab6201 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
<b>Dot blot</b>		Use a concentration of 1 - 10 µg/ml.
<b>ICC/IF</b>		Use a concentration of 5 µg/ml.
<b>ELISA</b>		Use a concentration of 0.1 µg/ml.
<b>Neutralising</b>		Use a concentration of 2 - 10 µg/ml.
<b>WB</b>	★★★★★ (1)	Use a concentration of 1 - 10 µg/ml. WB analysis of BDNF can give multiple bands, especially when using tissue lysates, because of many BDNF binding proteins that can form SDS-stable complexes even after boiling. Even in reduced samples, BDNF forms easily dimers and multimers, thus six or more bands can be present (most prominent: 28kDa dimer, monomer: ~14kDa). ab6201 will react also with BDNF-proforms
<b>IHC-FoFr</b>		Use a concentration of 1 - 5 µg/ml. PubMed: 18971468 Using ABC or tyramide amplification may be required to enhance the signal.

## Target

<b>Function</b>	During development, promotes the survival and differentiation of selected neuronal populations of the peripheral and central nervous systems. Participates in axonal growth, pathfinding and in the modulation of dendritic growth and morphology. Major regulator of synaptic transmission and plasticity at adult synapses in many regions of the CNS. The versatility of BDNF is emphasized by its contribution to a range of adaptive neuronal responses including long-term potentiation (LTP), long-term depression (LTD), certain forms of short-term synaptic plasticity, as well as homeostatic regulation of intrinsic neuronal excitability.
<b>Tissue specificity</b>	Brain. Highly expressed in hippocampus, amygdala, cerebral cortex and cerebellum. Also expressed in heart, lung, skeletal muscle, testis, prostate and placenta.
<b>Involvement in disease</b>	Bulimia nervosa 2 Congenital central hypoventilation syndrome
<b>Sequence similarities</b>	Belongs to the NGF-beta family.
<b>Post-translational modifications</b>	The propeptide is N-glycosylated and glycosulfated. Converted into mature BDNF by plasmin (PLG).
<b>Cellular localization</b>	Secreted.

## Images



**All lanes :** Anti-BDNF antibody (ab6201)

**Lane 1 :** Purified rHBNDF at 0.05  $\mu$ g

**Lane 2 :** Mouse brain lysate at 15  $\mu$ g

**Observed band size:** 14,28 kDa



ab6201 BDNF antibody immunostaining in the ventral horn of the spinal cord. Protocol: free floating, PFA/picric acid perfusion fixed rat coronal sections (30 microns) were incubated overnight in ab6201 (1/500) followed by 1 night of postfixation.

Immunofluorescence was visualised with TSA amplification.



ab6201 BDNF antibody immunostaining BDNF-containing hippocampal neurons. Protocol: PFA(only)-fixed free floating coronal rat brain sections (30 microns) were incubated for 3 days with ab6201 at 1/100 followed by direct immunofluorescence detection (alexa488; 1/1000). We recommend using an amplification IHC protocol following PFA/picric acid fixation of brain tissue for optimising BDNF detection using ab6201

Immunocytochemistry/ Immunofluorescence - Anti-  
BDNF antibody (ab6201)

ICC/IF image of ab6201 stained PC12 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab6201, 5µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Western blot - Anti-BDNF antibody (ab6201)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors