


Product datasheet

Anti-PSKH1 antibody ab69993

1 Image

Overview

Product name	Anti-PSKH1 antibody
Description	Rabbit polyclonal to PSKH1
Host species	Rabbit
Specificity	Detects endogenous levels of total PSKH1 protein.
Tested applications	Suitable for: WB, ELISA
Species reactivity	Reacts with: Human Predicted to work with: Mouse 
Immunogen	Synthesized peptide derived from an internal sequence of human PSKH1.
Positive control	Extracts from A549 cells

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS, 150mM Sodium chloride, pH 7.4
Purity	Immunogen affinity purified
Purification notes	Affinity purified from rabbit antiserum by affinity chromatography, using epitope specific immunogen.
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab69993** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		1/500 - 1/1000. Detects a band of approximately 48 kDa (predicted molecular weight: 48 kDa).

Application	Abreviews	Notes
ELISA		1/20000.

Target

Function	May be a SFC-associated serine kinase (splicing factor compartment-associated serine kinase) with a role in intranuclear SR protein (non-snRNP splicing factors containing a serine/arginine-rich domain) trafficking and pre-mRNA processing.
Tissue specificity	Expressed in all tissues and cell lines tested with the highest level of abundance in testis.
Sequence similarities	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. Contains 1 protein kinase domain.
Post-translational modifications	Autophosphorylated on serine residues. Myristoylated. Required for membrane association. Prerequisite for palmitoylation to occur. Palmitoylated.
Cellular localization	Golgi apparatus. Cytoplasm > cytoskeleton > centrosome. Nucleus speckle. Endoplasmic reticulum membrane. Cell membrane. Cytoplasm. Localized in the brefeldin A-sensitive Golgi compartment, at centrosomes, in the nucleus with a somewhat speckle-like presence, membrane-associated to the endoplasmic reticulum (ER) and the plasma membrane (PM), and more diffusely in the cytoplasm. Found to concentrate in splicing factor compartments (SFCs) within the nucleus of interphase cells. The acylation-negative form may be only cytoplasmic and nuclear. Acylation seems to allow the sequestering to the intracellular membranes. Myristoylation may mediate targeting to the intracellular non-Golgi membranes and palmitoylation may mediate the targeting to the Golgi membranes. Dual acylation is required to stabilize the interaction with Golgi membranes.

Images



All lanes : Anti-PSKH1 antibody (ab69993) at 1/500 dilution

Lane 1 : Extracts from A549 cells, minus immunising peptide

Lane 2 : Extracts from A549 cells, plus immunising peptide at 5µg

Lysates/proteins at 5 µg per lane.

Predicted band size: 48 kDa

Observed band size: 48 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors