


Product datasheet

Anti-ID2 antibody ab87189

[1 Abreviews](#)
[1 References](#)
[1 Image](#)

Overview

Product name	Anti-ID2 antibody
Description	Rabbit polyclonal to ID2
Host species	Rabbit
Tested applications	Suitable for: WB
Species reactivity	Reacts with: Mouse, Human Predicted to work with: Rat, Cow, Pig 
Immunogen	Synthetic peptide corresponding to a region within internal sequence amino acids 85-134 (IVSLHHQRPG QNQASRTPLT TLNTDISILS LQASEFPSEL MSNDSKALCG) of human ID2 (NP_002157). Run BLAST with ExPASy Run BLAST with NCBI
Positive control	Human fetal thymus nuclear lysate

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
Storage buffer	Preservative: 0.09% Sodium azide Constituents: PBS, 2% Sucrose
Purity	Immunogen affinity purified
Purification notes	Purified by peptide affinity chromatography method.
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab87189** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes

Application	Abreviews	Notes
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 15 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

Target

Function	ID (inhibitor of DNA binding) HLH proteins lack a basic DNA-binding domain but are able to form heterodimers with other HLH proteins, thereby inhibiting DNA binding. ID-2 may be an inhibitor of tissue-specific gene expression.
Tissue specificity	Highly expressed in early fetal tissues, including those of the central nervous system.
Sequence similarities	Contains 1 basic helix-loop-helix (bHLH) domain.
Developmental stage	Found in most early fetal tissues but not in the corresponding mature tissues.
Cellular localization	Cytoplasm. Nucleus.

Images



Anti-ID2 antibody (ab87189) at 1 µg/ml + human fetal thymus lysate at 10 µg

Secondary

HRP conjugated anti-Rabbit IgG at 1/50000 dilution

Predicted band size: 15 kDa

Observed band size: 15 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors