


Product datasheet

Anandamide (ethanol solution), endogenous cannabinoid ab120087

3 References 1 Image

Overview


| | |
|-------------------------------|---|
| Product name | Anandamide (ethanol solution), endogenous cannabinoid |
| Description | Endogenous cannabinoid |
| Biological description | Endogenous cannabinoid, binding to CB ₁ , CB ₂ and vanilloid receptors. Cannabimimetic <i>in vivo</i> . |
| Purity | > 98% |
| CAS Number | 94421-68-8 |
| Chemical structure | |



Properties

| | |
|-----------------------------|---|
| Chemical name | (5Z,8Z,11Z,14Z)-N-(2-Hydroxyethyl)icosa-5,8,11,14-tetraenamide |
| Molecular weight | 347.54 |
| Molecular formula | C ₂₂ H ₃₇ NO ₂ |
| PubChem identifier | 5281969 |
| Storage instructions | Store at -20°C (desiccating conditions). |
| Solubility overview | Supplied in ethanol (5 mg/ml) |
| Handling | <p>Providing storage is as stated on the product vial and the vial is kept tightly sealed, the product can be stored for up to 6 months. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Need more advice on solubility, usage and handling? Please visit our frequently asked questions (FAQ) page for more details.</p> |
| SMILES | O=C(CCC/C=CC/C=CC/C=CC/C=CCCC)NCCO |
| Source | Synthetic |

Images



Functional Studies - Anandamide (ethanol solution), endogenous cannabinoid (ab120087)

ab81298 staining FAK (phospho Y397) in SK-N-SH cells treated with anandamide (ethanol solution) (ab120087), by ICC/IF. Increase in FAK (phospho Y397) expression correlates with increased concentration of anandamide (ethanol solution), as described in literature.

The cells were incubated at 37°C for 10 minutes in media containing different concentrations of ab120087 (anandamide (ethanol solution)) in ethanol, fixed with 100% methanol for 5 minutes at -20°C and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with **ab81298** (5 μg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue. Membranes are stained in red with WGA.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team