

Product datasheet

Anti-ATG16L1 (phospho S278) antibody [EPR19016] ab195242

Recombinant RabMAb

1 References 6 Images

Overview

Product name	Anti-ATG16L1 (phospho S278) antibody [EPR19016]
Description	Rabbit monoclonal [EPR19016] to ATG16L1 (phospho S278)
Host species	Rabbit
Tested applications	Suitable for: ICC/IF, IHC-P, Dot blot, WB
Species reactivity	Reacts with: Mouse
Immunogen	<p>This product was produced with the following immunogens:</p> <p>Synthetic peptide within Mouse ATG16L1 aa 250-350 (phospho S278). The exact sequence is proprietary. Database link: Q8C0J2</p> <p>Synthetic peptide within Mouse ATG16L1 aa 250-350 (phospho S278). The exact sequence is proprietary. Database link: Q8C0J2</p>
Positive control	WB: HEK-293 overexpressing ATG16L1 (WT) whole cell lysate.
General notes	<p>Co-immunization performed with both peptides, clone obtained by screening with peptide 1.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	<p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 0.05% BSA, 40% Glycerol, PBS</p>
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR19016

Isotype

IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab195242** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/150.
IHC-P		1/300.
Dot blot		1/1000.
WB		Use at an assay dependent concentration. Predicted molecular weight: 68 kDa. For optimal WB signal, we recommend using 10X Blocking Buffer (ab126587).

Target

Function

Plays an essential role in autophagy: interacts with ATG12-ATG5 to mediate the conjugation of phosphatidylethanolamine (PE) to LC3 (MAP1LC3A, MAP1LC3B or MAP1LC3C), to produce a membrane-bound activated form of LC3 named LC3-II. Thereby, controls the elongation of the nascent autophagosomal membrane.

Involvement in disease

Inflammatory bowel disease 10

Sequence similarities

Belongs to the WD repeat ATG16 family.
Contains 7 WD repeats.

Post-translational modifications

Proteolytic cleavage by activated CASP3 leads to degradation and may regulate autophagy upon cellular stress and apoptotic stimuli.


Cellular localization

Cytoplasm. Preautophagosomal structure membrane. Recruited to omegasomes membranes by WIPI2. Omegasomes are endoplasmic reticulum connected structures at the origin of preautophagosomal structures. Localized to preautophagosomal structure (PAS) where it is involved in the membrane targeting of ATG5. Localizes also to discrete punctae along the ciliary axoneme.

Form

There are 4 isoforms produced by alternative splicing.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr. Ryan Russell (University of Ottawa).

IHC images of mice quadriceps showing either pATG16L1 or LC3B staining

Mice were fed ad libitum or starved for 16 hours. Quadriceps muscle were immediately harvested and fixed in 10% formalin for 2 days. The samples were then paraffin embedded, sectioned into 4µm thick slices, and mounted onto glass microscope slides. Slides were stained with primary antibody overnight at 4°C: LC3B 1/1000, pATG16L1 (ab195242) 1/300. Secondary antibody: Alexa Fluor 555 anti-rabbit, 1/1000.




Immunocytochemistry/ Immunofluorescence - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr. Ryan Russell (University of Ottawa).

IF showing pATG16L1 (red) and total ATG16L1 (green):

Polyclonal population of ATG16L1 KO and HA-ATG16L1 reconstituted cells were starved of amino acid for 1 hour and stained. Blocking buffer used for pATG16L1 staining: 0.1% BSA, 1x abcam blocking buffer ab126587, diluted in PBS. Anti-pATG16L1 (ab195242) concentration: 1/150. Anti-ATG16L1, concentration: 1/200 Secondary antibody (Alexa Fluor 647/488) concentration: 1/1000.




Immunocytochemistry/ Immunofluorescence - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr. Ryan Russell (University of Ottawa).

IF showing pATG16L1 (red), LC3B (green) and p62 (white):


MEF cells were amino acid starved for 1 hour. Blocking buffer used for pATG16L1 (ab195242): 0.1% BSA, 1x abcam blocking buffer (ab126587), diluted in PBS. Anti-pATG16L1 (ab195242) concentration: 1/150. Secondary antibody (Alexa Fluor 647) concentration: 1/1000



Western blot - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr Ryan Russell (University of Ottawa).

HCT116 wild-type and ATG16L1 knockout cells were incubated with either complete media or amino acid deficient DMEM for 3 hours. 5ug of whole cell lysate were resolved by SDS-PAGE on a 6%-18% gradient gel, then transferred onto PVDF membrane. Membrane was blocked in 10X blocking buffer (Cat # [ab126587](#)) diluted in TBS solution for 30 minutes; incubated with 1:1000 primary antibody in 2.5% BSA TBST solution overnight at 4°C ; incubated with 1:15000 secondary antibody in 2% milk TBST solution for 45 minutes. Immobilon ECL was applied for 1 minute then imaged with film.



Western blot - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

All lanes : Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242) at 1/1000 dilution

Lane 1 : HEK-293 (human epithelial cell line from embryonic kidney) transfected with an empty vector (vector control), containing a myc-His-tag®, whole cell lysate

Lane 2 : HEK-293 (human epithelial cell line from embryonic kidney) transfected with ATG16L1 (WT) expression vector containing a myc-His-tag®, whole cell lysate

Lane 3 : HEK-293 (human epithelial cell line from embryonic kidney) transfected with ATG16L1 (WT) expression vector containing a myc-His-tag®, followed by treatment with alkaline phosphatase for 1 hour, whole cell lysate

Lane 4 : HEK-293 (human epithelial cell line from embryonic kidney) transfected with ATG16L1 S278A expression vector containing a myc-His-tag®, whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Developed using the ECL technique.

Predicted band size: 68 kDa

Observed band size: 68 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDm/TBST.



Dot blot analysis of ATG16L1 (phospho S278) labeled with ab195242 at 1/1,000 dilution.

Lane 1: ATG16L1 (phospho S278) peptide;

Lane 2: ATG16L1 non-phospho peptide;

Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100,000 dilution was used as secondary antibody.

Blocking/Dilution buffer: 5% NFDm/TBST.

Exposure time: 3 minutes.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors