

## Product datasheet

# Anti-ATP1A3 antibody [XVIF9-G10] ab2826

★★★★★ 5 Abreviews 8 References 8 Images

### Overview

<b>Product name</b>	Anti-ATP1A3 antibody [XVIF9-G10]
<b>Description</b>	Mouse monoclonal [XVIF9-G10] to ATP1A3
<b>Host species</b>	Mouse
<b>Specificity</b>	The immunogen used for this product shares 89% homology with ATP1A2. Cross-reactivity with this protein has not been confirmed experimentally
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt, IHC-Fr, ICC/IF, WB, Inhibition Assay, IP, IHC-P
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Sheep, Rabbit, Guinea pig, Cow, Dog, Human, Pig, Shark <b>Predicted to work with:</b> Non human primates, Amphibian 
<b>Immunogen</b>	Full length native protein (purified) corresponding to Dog ATP1A3. Canine cardiac microsomes.
<b>General notes</b>	This product was previously labelled as alpha 3 Sodium Potassium ATPase

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	pH: 7.2 Preservative: 0.05% Sodium azide Constituents: 0.42% Potassium phosphate, 0.88% Sodium chloride
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	XVIF9-G10
<b>Isotype</b>	IgG1

### Applications

Our [Abpromise guarantee](#) covers the use of **ab2826** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use 1µg for 10 <sup>6</sup> cells. <a href="#">ab170190</a> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
IHC-Fr		Use a concentration of 3 µg/ml.
ICC/IF		Use a concentration of 5 µg/ml.
WB	★★★★☆	Use a concentration of 1 µg/ml.
Inhibition Assay		1/10.
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.

## Target

<b>Function</b>	This is the catalytic component of the active enzyme, which catalyzes the hydrolysis of ATP coupled with the exchange of sodium and potassium ions across the plasma membrane. This action creates the electrochemical gradient of sodium and potassium ions, providing the energy for active transport of various nutrients.
<b>Involvement in disease</b>	Dystonia 12 Alternating hemiplegia of childhood 2 Cerebellar ataxia, areflexia, pes cavus, optic atrophy, and sensorineural hearing loss
<b>Sequence similarities</b>	Belongs to the cation transport ATPase (P-type) (TC 3.A.3) family. Type IIC subfamily.
<b>Cellular localization</b>	Cell membrane.

## Images




Western blot - Anti-ATP1A3 antibody [XVIF9-G10] (ab2826)

This image is courtesy of an anonymous Abreview

Anti-ATP1A3 antibody [XVIF9-G10] (ab2826) at 1/1000 dilution (5% Milk for 1 hour 30 minutes at 24°C.) + Human brain cortex whole tissue lysate (10µg). with 5% Milk for 1 hour for 24°C.

### Secondary


An HRP-conjugated goat anti-mouse IgG1 at 1/5000 dilution



Flow Cytometry - Anti-ATP1A3 antibody [XVIF9-G10] (ab2826)


Overlay histogram showing SH-SY5Y cells stained with ab2826 (red line). The cells were fixed with 4% paraformaldehyde and incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab2826, 1µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (ab96879) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (ab91353, 2µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed.

Please note that Abcam do not have any data for use of this antibody on non-fixed cells. We welcome any customer feedback.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATP1A3 antibody [XVIF9-G10] (ab2826)

Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized Human prostate carcinoma tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:50 with a mouse monoclonal antibody recognizing Sodium/Potassium ATPase alpha-3 ab2826 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATP1A3 antibody [XVIF9-G10] (ab2826)

Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized Human colon tissue tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:200 with a mouse monoclonal antibody recognizing Sodium/Potassium ATPase alpha-3 ab2826 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATP1A3 antibody [XVIF9-G10] (ab2826)

Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized Human tonsil tissue tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:200 with a mouse monoclonal antibody recognizing Sodium/Potassium ATPase alpha-3 ab2826 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.




Immunocytochemistry/ Immunofluorescence - Anti-ATP1A3 antibody [XVIF9-G10] (ab2826)

Immunofluorescent analysis of Sodium/Potassium ATPase alpha-3 using Sodium/Potassium ATPase alpha-3 Monoclonal antibody (XVIF9-G10) ab2826 shows staining in C6 glioma cells. Sodium/Potassium ATPase alpha-3 staining (green) F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Sodium/Potassium ATPase alpha-3 ab2826 at a dilution of 1:20 over night at 4 °C washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-ATP1A3 antibody [XVIF9-G10] (ab2826)

Immunofluorescent analysis of Sodium/Potassium ATPase alpha-3 using Sodium/Potassium ATPase alpha-3 Monoclonal antibody (XVIF9-G10) ab2826 shows staining in U251 glioma cells. Sodium/Potassium ATPase alpha-3 staining (green) F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Sodium/Potassium ATPase alpha-3 ab2826 at a dilution of 1:20 over night at 4 °C washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-ATP1A3 antibody [XVIF9-G10] (ab2826)

Immunofluorescent analysis of Sodium/Potassium ATPase alpha-3 using Sodium/Potassium ATPase alpha-3 Monoclonal antibody (XVIF9-G10) ab2826 shows staining in HeLa cells. Sodium/Potassium ATPase alpha-3 staining (green) F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Sodium/Potassium ATPase alpha-3 ab2826 at a dilution of 1:20 over night at 4 °C washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors