

## Product datasheet

# Anti-beta Dystroglycan (phospho Y892) antibody [M117] ab29037

[2 Images](#)

### Overview

<b>Product name</b>	Anti-beta Dystroglycan (phospho Y892) antibody [M117]
<b>Description</b>	Mouse monoclonal [M117] to beta Dystroglycan (phospho Y892)
<b>Host species</b>	Mouse
<b>Specificity</b>	Ab29037 recognises beta Dystroglycan (phospho Y892).
<b>Tested applications</b>	<b>Suitable for:</b> WB, ELISA
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	Synthetic peptide conjugated to KLH corresponding to amino acid residues around tyrosine 892 of human dystroglycan.
<b>Positive control</b>	HepG2 cells or HeLa cells treated with pervanadate.

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
<b>Storage buffer</b>	Preservative: 0.05% Sodium Azide. Constituents: 50% Glycerol, PBS, 1mg/ml BSA.
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	M117
<b>Isotype</b>	IgG

### Applications

Our [Abpromise guarantee](#) covers the use of **ab29037** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		

Application	Abreviews	Notes
-------------	-----------	-------

ELISA

**Application notes**

ELISA: 1/2000.  
 WB: 1/1000. Membrane was incubated with diluted antibody in 5% non fat milk, PBS, 0.04% Tween 20 for 1 hour at room temperature. Predicted molecular weight: 97 kDa (Dystroglycan is expressed as a precursor protein that is cleaved into alpha and beta dystroglycan. The beta dystroglycan fragment is about 43 kDa).

Not yet tested in other applications.  
 Optimal dilutions/concentrations should be determined by the end user.

**Target**


**Relevance**

Dystroglycans are essential elements of the neuromuscular junction (NMJ). The gene for dystroglycan is expressed as a precursor protein that is post translationally cleaved into a 156 kDa extracellular peripheral membrane protein called alpha dystroglycan and a 43 kDa transmembrane protein, beta dystroglycan. The latter protein contains a PPxY motif that promotes binding to WW domain containing proteins, such as utrophin and dystrophin. Phosphorylation at tyrosine 892 within the PPxY motif may regulate c Src interactions with beta dystroglycan, as well as inhibit interactions with WW domain proteins. In skeletal muscle, beta dystroglycan is normally localized to the plasma membrane, however phosphorylation of Tyr892 leads to localization of beta dystroglycan to endosomal compartments along with c Src. Thus, phosphorylation at Tyr892 may have important roles in altering the localization of beta dystroglycan during NMJ formation.

**Cellular localization**

Type I membrane protein.

**Images**



Western blot - Anti-beta Dystroglycan (phospho Y892) antibody [M117] (ab29037)

**All lanes :** Anti-beta Dystroglycan (phospho Y892) antibody [M117] (ab29037)

**Lane 1 :** Lysates from HepG2 cells (untreated)


**Lane 2 :** Lysates from HepG2 cells treated with pervanadate (1 mM) for 30 min

Lysates/proteins at 20 µg per lane.

**Predicted band size:** 97 kDa

**Observed band size:** 43 kDa

Dystroglycan is expressed as a precursor protein (97kDa) that is cleaved into alpha and beta dystroglycan. The beta dystroglycan fragment is about 43 kDa.



Western blot - Anti-beta Dystroglycan (phospho Y892) antibody [M117] (ab29037)

Western blot analysis of HepG2 cells untreated (lanes 1 & 3) or treated with pervanadate (1 mM) for 30 min (lanes 2 & 4). Blots were probed with anti-beta-dystroglycan (lanes 1 & 2) and anti-beta-dystroglycan (Tyr-892) (lanes 3 & 4).

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors