

Product datasheet

Anti-Caspase-8 antibody ab4052

★★★★★ 1 Abreviews 10 References 2 Images

Overview

Product name	Anti-Caspase-8 antibody
Description	Rabbit polyclonal to Caspase-8
Host species	Rabbit
Tested applications	Suitable for: IHC-P, IHC-Fr
Species reactivity	Reacts with: Mouse, Rat, Cow, Human, Monkey
Immunogen	Recombinant full length protein corresponding to Human Caspase-8. Database link: Q14790
Positive control	Stomach.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.05% Sodium azide Constituent: 1% BSA
Purity	IgG fraction
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab4052** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P	★★★★★	1/50 - 1/100. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
IHC-Fr		1/50 - 1/100.

Target

Function

Most upstream protease of the activation cascade of caspases responsible for the TNFRSF6/FAS mediated and TNFRSF1A induced cell death. Binding to the adapter molecule FADD recruits it to either receptor. The resulting aggregate called death-inducing signaling complex (DISC) performs CASP8 proteolytic activation. The active dimeric enzyme is then liberated from the DISC and free to activate downstream apoptotic proteases. Proteolytic fragments of the N-terminal propeptide (termed CAP3, CAP5 and CAP6) are likely retained in the DISC. Cleaves and activates CASP3, CASP4, CASP6, CASP7, CASP9 and CASP10. May participate in the GZMB apoptotic pathways. Cleaves ADPRT. Hydrolyzes the small-molecule substrate, Ac-Asp-Glu-Val-Asp-AMC. Likely target for the cowpox virus CRMA death inhibitory protein. Isoform 5, isoform 6, isoform 7 and isoform 8 lack the catalytic site and may interfere with the pro-apoptotic activity of the complex.

Tissue specificity

Isoform 1, isoform 5 and isoform 7 are expressed in a wide variety of tissues. Highest expression in peripheral blood leukocytes, spleen, thymus and liver. Barely detectable in brain, testis and skeletal muscle.

Involvement in disease

Defects in CASP8 are the cause of caspase-8 deficiency (CASP8D) [MIM:607271]. CASP8D is a disorder resembling autoimmune lymphoproliferative syndrome (ALPS). It is characterized by lymphadenopathy, splenomegaly, and defective CD95-induced apoptosis of peripheral blood lymphocytes (PBLs). It leads to defects in activation of T-lymphocytes, B-lymphocytes, and natural killer cells leading to immunodeficiency characterized by recurrent sinopulmonary and herpes simplex virus infections and poor responses to immunization.

Sequence similarities

Belongs to the peptidase C14A family.
Contains 2 DED (death effector) domains.

Domain

Isoform 9 contains a N-terminal extension that is required for interaction with the BCAP31 complex.

Post-translational modifications

Generation of the subunits requires association with the death-inducing signaling complex (DISC), whereas additional processing is likely due to the autocatalytic activity of the activated protease. GZMB and CASP10 can be involved in these processing events.
Phosphorylated upon DNA damage, probably by ATM or ATR.

Cellular localization

Cytoplasm.


Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Caspase-8 antibody (ab4052)

This image is courtesy of an Abreview submitted by Dr B Goilav

ab4052 staining Caspase-8 in human kidney tissue section by immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Tissue underwent fixation in paraformaldehyde, heat mediated antigen retrieval in 10mM Citrate buffer pH 6.0, permeabilization in 0.1% Triton X-100 prior to blocking in 10% serum for 20 minutes at 22°C. The primary antibody, diluted 1/50 in PBS with 2% BSA and incubated with sample, for 45 minutes at 22°C. An Avidin-peroxidase conjugated horse polyclonal to rabbit Ig, was used undiluted as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Caspase-8 antibody (ab4052)

ab4052 staining Caspase-8 in formalin-fixed, paraffin-embedded Human stomach tissue by Immunohistochemistry.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors