abcam

Product datasheet

PE Anti-CD19 antibody [1G9] ab52056

1 Image

Overview

Product name	PE Anti-CD19 antibody [1G9]
Description	PE Mouse monoclonal [1G9] to CD19
Host species	Mouse
Conjugation	PE. Ex: 488nm, Em: 575nm
Tested applications	**Suitable for:** Flow Cyt
Species reactivity	**Reacts with:** Human
Immunogen	The details of the immunogen for this antibody are not available.
Positive control	Peripheral blood lymphocytes
General notes	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.08% Sodium azide Constituent: PBS 2% protein carrier.
Clonality	Monoclonal
Clone number	1G9
Isotype	IgG1
Light chain type	kappa

Applications

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab52056 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use 10µl for 10^6 cells. in 100µl, or for 100µl of whole blood.

Target

Function Assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation.

Involvement in disease Defects in CD19 are the cause of immunodeficiency common variable type 3 (CVID3) [MIM:613493]; also called antibody deficiency due to CD19 defect. CVID3 is a primary immunodeficiency characterized by antibody deficiency, hypogammaglobulinemia, recurrent bacterial infections and an inability to mount an antibody response to antigen. The defect results from a failure of B-cell differentiation and impaired secretion of immunoglobulins; the numbers of circulating B cells is usually in the normal range, but can be low.

Sequence similarities Contains 2 Ig-like C2-type (immunoglobulin-like) domains.

Post-translational modifications Phosphorylated on serine and threonine upon DNA damage, probably by ATM or ATR. Phosphorylated on tyrosine following B-cell activation.

Cellular localization Membrane.

Images



Flow Cytometry - PE Anti-CD19 antibody [1G9] (ab52056)

FACS analysis of CD19 expression in Peripheral blood lymphocytes using ab52056.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit **https://www.abcam.com/abpromise** or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors