

Product datasheet

Anti-CD31 antibody [JC/70A] - BSA and Azide free ab212712

★★★★☆ 1 Abreviews 1 References 4 Images

Overview

Product name	Anti-CD31 antibody [JC/70A] - BSA and Azide free
Description	Mouse monoclonal [JC/70A] to CD31 - BSA and Azide free
Host species	Mouse
Tested applications	Suitable for: IHC-P, WB, Flow Cyt
Species reactivity	Reacts with: Human Predicted to work with: Rabbit, Cynomolgus monkey Does not react with: Rat, Pig
Immunogen	Tissue, cells or virus corresponding to Human CD31. Membrane preparation of a spleen from a patient with hairy cell leukemia. Database link: P16284
Positive control	IHC-P: Human colon carcinoma and Human Angiosarcoma tissues; Flow Cyt: Jurkat cells WB: THP-1 cell lysates
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Constituent: 100% PBS
Carrier free	Yes
Purity	Protein A/G purified
Clonality	Monoclonal

Clone number	JC/70A
Isotype	IgG1
Light chain type	kappa

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab212712 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P	★★★★★ (1)	<p>Use a concentration of 1 - 2 µg/ml. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.</p> <p>The ideal fixation time will depend on the size of the tissue block and the type of tissue, but fixation between 18–24h is suitable for most samples.</p> <p>Positive Control: HU tonsil tissue</p>
WB		<p>Use a concentration of 0.5 - 1 µg/ml. Predicted molecular weight: 82 kDa.</p> <p>Treat samples with PNGase F or phosphatase to confirm the specificity of bands if necessary.</p> <p>The observed band size of CD31 may not be the same as predicted MWs in WB due to the different forms and modifications of CD31.</p>
Flow Cyt		Use 0.5-1µg for 10 ⁶ cells.

Target

Function

Induces susceptibility to atherosclerosis (By similarity). Cell adhesion molecule which is required for leukocyte transendothelial migration (TEM) under most inflammatory conditions. Tyr-690 plays a critical role in TEM and is required for efficient trafficking of PECAM1 to and from the lateral border recycling compartment (LBRC) and is also essential for the LBRC membrane to be targeted around migrating leukocytes. Prevents phagocyte ingestion of closely apposed viable cells by transmitting 'detachment' signals, and changes function on apoptosis, promoting tethering of dying cells to phagocytes (the encounter of a viable cell with a phagocyte via the homophilic interaction of PECAM1 on both cell surfaces leads to the viable cell's active repulsion from the phagocyte. During apoptosis, the inside-out signaling of PECAM1 is somehow disabled so that the apoptotic cell does not actively reject the phagocyte anymore. The lack of this repulsion signal together with the interaction of the eat-me signals and their respective receptors causes the attachment of the apoptotic cell to the phagocyte, thus triggering the process of engulfment). Isoform Delta15 is unable to protect against apoptosis. Modulates BDKRB2 activation. Regulates bradykinin- and hyperosmotic shock-induced ERK1/2 activation in human umbilical cord vein cells (HUVEC).

Tissue specificity

Expressed on platelets and leukocytes and is primarily concentrated at the borders between endothelial cells. Isoform Long predominates in all tissues examined. Isoform Delta12 is detected only in trachea. Isoform Delta14-15 is only detected in lung. Isoform Delta14 is detected in all tissues examined with the strongest expression in heart. Isoform Delta15 is expressed in brain,

testis, ovary, cell surface of platelets, human umbilical vein endothelial cells (HUVECs), Jurkat T-cell leukemia, human erythroleukemia (HEL) and U937 histiocytic lymphoma cell lines (at protein level).

Sequence similarities

Contains 6 Ig-like C2-type (immunoglobulin-like) domains.

Domain

The Ig-like C2-type domains 2 and 3 contribute to formation of the complex with BDKRB2 and in regulation of its activity.


Post-translational modifications

Phosphorylated on Ser and Tyr residues after cellular activation. In endothelial cells Fyn mediates mechanical-force (stretch or pull) induced tyrosine phosphorylation.

Cellular localization

Membrane. Cell junction. Localizes to the lateral border recycling compartment (LBRC) and recycles from the LBRC to the junction in resting endothelial cells and Cell junction. Localizes to the lateral border recycling compartment (LBRC) and recycles from the LBRC to the junction in resting endothelial cells.


Images



Anti-CD31 antibody [JC/70A] - BSA and Azide free (ab212712) + THP-1 cell lysate


Predicted band size: 82 kDa

Western blot - Anti-CD31 antibody [JC/70A] - BSA and Azide free (ab212712)




Flow Cytometry analysis of Jurkat cells labeling CD31 with ab212712 (Green). Black: cells alone; Grey: Isotype Control.

Flow Cytometry - Anti-CD31 antibody [JC/70A] - BSA and Azide free (ab212712)



Immunohistochemical analysis of formalin-fixed, paraffin-embedded Human colon carcinoma tissue labeling CD31 with ab212712 at 2 µg/ml.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD31 antibody [JC/70A] - BSA and Azide free (ab212712)



Immunohistochemical analysis of formalin-fixed, paraffin-embedded Human angiosarcoma tissue labeling CD31 with ab212712 at 2 µg/ml dilution.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD31 antibody [JC/70A] - BSA and Azide free (ab212712)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors