

## Product datasheet

# Anti-CD68 antibody [KP1] - BSA and Azide free ab233172

Recombinant

★★★★☆ 1 Abreviews 6 Images

### Overview

<b>Product name</b>	Anti-CD68 antibody [KP1] - BSA and Azide free
<b>Description</b>	Mouse monoclonal [KP1] to CD68 - BSA and Azide free
<b>Host species</b>	Mouse
<b>Tested applications</b>	<b>Suitable for:</b> Mass Cytometry, ICC/IF, IHC-P, WB <b>Unsuitable for:</b> Flow Cyt
<b>Species reactivity</b>	<b>Reacts with:</b> Human
<b>Immunogen</b>	Tissue, cells or virus. This information is considered to be commercially sensitive.
<b>Positive control</b>	WB: Human spleen lysate. IHC-P: Human tonsil and liver tissue. ICC/IF: THP-1 cells. IMC: Human tonsil tissue
<b>General notes</b>	ab233172 is the carrier-free version of <a href="#">ab955</a> .

Our [carrier-free](#) antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our [conjugation kits](#) for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C. Do Not Freeze.
<b>Storage buffer</b>	pH: 7.2

	Constituent: PBS
<b>Carrier free</b>	Yes
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	KP1
<b>Isotype</b>	IgG1
<b>Light chain type</b>	kappa

## Applications

**The Abpromise guarantee** Our [Abpromise guarantee](#) covers the use of ab233172 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
<b>Mass Cytometry</b>	★★★★★ (1)	Use at an assay dependent concentration.
<b>ICC/IF</b>		1/50.
<b>IHC-P</b>		1/3000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
<b>WB</b>		1/1000. Detects a band of approximately 110 kDa (predicted molecular weight: 37 kDa).

**Application notes** Is unsuitable for Flow Cyt.

## Target

<b>Function</b>	Could play a role in phagocytic activities of tissue macrophages, both in intracellular lysosomal metabolism and extracellular cell-cell and cell-pathogen interactions. Binds to tissue- and organ-specific lectins or selectins, allowing homing of macrophage subsets to particular sites. Rapid recirculation of CD68 from endosomes and lysosomes to the plasma membrane may allow macrophages to crawl over selectin-bearing substrates or other cells.
<b>Tissue specificity</b>	Highly expressed by blood monocytes and tissue macrophages. Also expressed in lymphocytes, fibroblasts and endothelial cells. Expressed in many tumor cell lines which could allow them to attach to selectins on vascular endothelium, facilitating their dissemination to secondary sites.
<b>Sequence similarities</b>	Belongs to the LAMP family.
<b>Post-translational modifications</b>	N- and O-glycosylated.
<b>Cellular localization</b>	Cell membrane and Endosome membrane. Lysosome membrane.

## Images




Mass Cytometry - Anti-CD68 antibody [KP1] - BSA and Azide free (ab233172)

This image is courtesy of the Single Cell & Imaging Mass Cytometry Analysis Platform, Goodman Cancer Research Centre, McGill University

**Imaging Mass Cytometry™ (IMC™)** image of human tonsil tissue stained with Anti-CD68 antibody [KP1]. ab233172 (carrier-free antibody, purified) was metal-conjugated using a Maxpar® Antibody Labeling Kit from Fluidigm. Immunostaining was performed according to Fluidigm's protocols. Briefly, slides were subject to deparaffinization and heat-induced epitope retrieval, followed by overnight incubation at 4°C with an antibody cocktail containing metal-tagged antibodies in blocking buffer. Slides were subsequently washed with 0.2% Triton-X and 1x PBS, counterstained with Cell-ID™ Intercalator-Ir diluted at 1/400 in 1x PBS for 30 min at room temperature, rinsed for 5 min with distilled H<sub>2</sub>O, and air-dried prior to IMC™ acquisition. IMC™ acquisition was performed using the Fluidigm Hyperion™ Imaging System.

*Imaging Mass Cytometry™, IMC™, Cell-ID™, Hyperion™ and Maxpar® are trademarks of Fluidigm Canada*




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD68 antibody [KP1] - BSA and Azide free (ab233172)

Immunohistochemical analysis of paraffin-embedded human liver tissue labeling CD68 with ab233172 at 1/3000 dilution, followed by Goat Anti-mouse IgG H&L (HRP) ready to use. Cytoplasmic staining on Kupffer cells of human liver (PMID: 12118106) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-mouse IgG H&L (HRP) ready to use.

Heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD68 antibody [KP1] - BSA and Azide free (ab233172)

Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling CD68 with ab233172 at 1/3000 dilution, followed by Goat Anti-mouse IgG H&L (HRP) ready to use. Cytoplasmic staining on macrophages of human tonsil (PMID: 19543531) is observed. Counter stained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-mouse IgG H&L (HRP) ready to use.


Heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0).



Immunocytochemistry/ Immunofluorescence - Anti-CD68 antibody [KP1] - BSA and Azide free (ab233172)

Immunofluorescent analysis of 100% methanol-fixed THP-1 (human monocytic leukemia cell line) cells labeling CD68 with ab233172 at 1/50 dilution, followed by Goat Anti-Mouse IgG H&L (Alexa Fluor<sup>®</sup> 488) (ab150113) secondary antibody at 1/1000 dilution (green). Confocal image showing cytoplasmic staining in THP-1 cells. The nuclear counter stain is DAPI (blue). Tubulin is detected with Recombinant Anti-beta IV Tubulin antibody [EPR16775] (ab179504) at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor<sup>®</sup> 594) (ab150080) secondary antibody at 1/1000 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, followed by Goat Anti-Mouse IgG H&L (Alexa Fluor<sup>®</sup> 488) (ab150113) secondary antibody at 1/1000 dilution.



Western blot - Anti-CD68 antibody [KP1] - BSA and Azide free (ab233172)

Anti-CD68 antibody [KP1] - BSA and Azide free (ab233172) at 1/1000 dilution + Human spleen lysate at 20 µg/ml

**Secondary**

Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/10000 dilution

**Predicted band size:** 37 kDa

**Observed band size:** 110 kDa

**Exposure time:** 70 seconds

Blocking and dilution buffer: 5% NFDM/TBST.

The observed molecular weight is consistent with the literature (PMID:18405323; PMID:11739566; PMID: 16710801).

Why choose a recombinant antibody?

 <p><b>Research with confidence</b> Consistent and reproducible results</p>	 <p><b>Long-term and scalable supply</b> Recombinant technology</p>
 <p><b>Success from the first experiment</b> Confirmed specificity</p>	 <p><b>Ethical standards compliant</b> Animal-free production</p>

Anti-CD68 antibody [KP1] - BSA and Azide free (ab233172)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Promise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors