

Product datasheet

Anti-CDKN2A/p14ARF antibody ab3642

14 References 3 Images

Overview

|                            |   |
|----------------------------|---|
| <b>Product name</b>        | Anti-CDKN2A/p14ARF antibody   |
| <b>Description</b>         | Rabbit polyclonal to CDKN2A/p14ARF  |
| <b>Host species</b>        | Rabbit  |
| <b>Tested applications</b> | <b>Suitable for:</b> IHC-P, IHC-Fr, WB, ICC/IF  |
| <b>Species reactivity</b>  | <b>Reacts with:</b> Human   |
| <b>Immunogen</b>           | Synthetic peptide within Human CDKN2A/p14ARF (C terminal). The exact sequence is proprietary. |
| <b>Positive control</b>    | ICC/IF: HeLa cells. IHC-P: Lymphoma tissue.   |

Properties

|                             |   |
|-----------------------------|---|
| <b>Form</b>                 | Liquid  |
| <b>Storage instructions</b> | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| <b>Storage buffer</b>       | pH: 7.6<br>Preservative: 0.1% Sodium azide<br>Constituents: PBS, 1% BSA               |
| <b>Purity</b>               | Immunogen affinity purified   |
| <b>Clonality</b>            | Polyclonal  |
| <b>Isotype</b>              | IgG   |

Applications

Our [Abpromise guarantee](#) covers the use of **ab3642** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes   |
|-------------|-----------|---|
| IHC-P       |           | 1/25. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. |
| IHC-Fr      |           | Use at an assay dependent concentration.  |

| Application | Abreviews | Notes  |
|-------------|-----------|--|
| WB          |           | Use a concentration of 1 - 2 µg/ml. Detects a band of approximately 14 kDa (predicted molecular weight: 14 kDa). |
| ICC/IF      |           | Use a concentration of 1 µg/ml.  |

## Target


### Relevance

The gene for CDKN2A generates several transcripts/proteins which differ from each other in their first exons. Three of these transcripts are generated by alternative splicing (isoform 1 a.k.a p16INK4A, isoform 2 and isoform 3 a.k.a p12), two of which are known to function as inhibitors of CDK4 kinase. One other transcript that is generated from this gene contains an alternate reading frame (ARF), with the first exon located 20kb upstream of the remainder of the gene (isoform 4 a.k.a. p14ARF, p19ARF, ARF). In spite of the structural and some functional differences, all the proteins encoded by the CDKN2A gene are involved in cell cycle G1 control.

### Cellular localization

Cytoplasmic and Nuclear


## Images



Confocal image showing nuclear staining increased after HeLa cells transfected with CDKN2A/p14ARF.


ICC/IF image of ab3642 stained HeLa cells. The cells were 4% paraformaldehyde fixed and then incubated in 0.1% trixonX-100 to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with primary antibody ab3642 at a dilution of 1/250. An AlexaFluor®594 goat anti-mouse secondary IgG (ab150077) was used at a 1/1000 dilution. Anti-tubulin (ab7291) and an AlexaFluor®594 goat anti-mouse IgG (ab150120) were used as counterstains, both at a dilution of 1/1000. DAPI was used to stain the cell nuclei blue.

Immunocytochemistry/ Immunofluorescence - Anti-CDKN2A/p14ARF antibody (ab3642)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CDKN2A/p14ARF antibody (ab3642)

This image was kindly supplied as part of the review submitted by Joost van Galen.



Immunocytochemistry/ Immunofluorescence - Anti-CDKN2A/p14ARF antibody (ab3642)

ICC/IF image of ab3642 stained HeLa cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab3642 at 1µg/ml overnight at +4°C. The secondary antibody (green) was DyLight® 488 goat anti- rabbit (ab96899) IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors