

Product datasheet

Anti-CDKN2A/p16INK4a antibody [EPR20418] ab211542

Recombinant RabMAb

★★★★★ 7 Abreviews 28 References 6 Images

Overview

Product name	Anti-CDKN2A/p16INK4a antibody [EPR20418]
Description	Rabbit monoclonal [EPR20418] to CDKN2A/p16INK4a
Host species	Rabbit
Specificity	Expression levels of the CDKN2A/p16INK4a protein may vary with sample type. It's barely expressed in normal tissue, and mostly expressed in some tumour tissues, such as cervical cancer, breast cancer and so on. Moreover, only expressed in some cell lines. Please see images for recommended positive controls.
Tested applications	Suitable for: Flow Cyt (Intra), WB, IP, ICC/IF
Species reactivity	Reacts with: Mouse
Immunogen	Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: MEF whole cell lysate; His-tagged mouse CDKN2A/p16INK4a recombinant protein, aa1-168. ICC/IF: MEF cells. Flow Cyt (intra): MEF cells. IP: MEF whole cell lysate.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 0.05% BSA, 40% Glycerol, PBS
Purity	Protein A purified

Clonality	Monoclonal
Clone number	EPR20418
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab211542 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
Flow Cyt (Intra)		1/500.
WB	★★★★★ (2)	1/2000. Detects a band of approximately 16, 14 kDa (predicted molecular weight: 18 kDa).
IP		1/30.
ICC/IF	★★★★★ (2)	1/100.

Target

Cellular localization Cytoplasmic and Nuclear

Form There are 4 isoforms produced by alternative splicing. Isoform 1 also known as: p16INK4a; Isoform 3 also known as: p12; Isoform 4 also known as: p14ARF; p19ARF; ARF.

Images



Immunofluorescent analysis of 100% methanol-fixed MEF (mouse embryonic fibroblast cell line) and NIH/3T3 (mouse embryo fibroblast cell line) cells labeling CDKN2A/p16INK4a with ab211542 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1,000 dilution (green). Confocal image showing nuclear and cytoplasmic staining on MEF cell line.


Negative control: NIH/3T3 (PMID: 15210712).

The nuclear counter stain is DAPI (blue). Tubulin is detected with Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) (ab195889) (red) at 1/200 dilution.


Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1,000 dilution.

MEF cells were kindly provided by professor Pinlong Xu, Zhejiang

University.



Flow Cytometry (Intracellular) - Anti-CDKN2A/p16INK4a antibody [EPR20418] (ab211542)



Western blot - Anti-CDKN2A/p16INK4a antibody [EPR20418] (ab211542)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed MEF (mouse embryonic fibroblast cell line) cell line labeling CDKN2A/p16INK4a with ab211542 at 1/500 dilution (red) compared with a Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) at 1/2000 dilution was used as the secondary antibody.

MEF cells were kindly provided by professor Pinlong Xu, Zhejiang University.

All lanes : Anti-CDKN2A/p16INK4a antibody [EPR20418] (ab211542) at 1/1000 dilution

Lane 1 : His-tagged mouse CDKN2A/p16INK4a recombinant protein

Lane 2 : His-tagged mouse CDKN2B/p15INK4b recombinant protein

Lysates/proteins at 0.01 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution


Developed using the ECL technique.

Predicted band size: 18 kDa

Observed band size: 18 kDa

Exposure time: 1 second

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-CDKN2A/p16INK4a antibody [EPR20418] (ab211542)

All lanes : Anti-CDKN2A/p16INK4a antibody [EPR20418] (ab211542) at 1/2000 dilution

Lane 1 : MEF (mouse embryonic fibroblast cell line) whole cell lysate

Lane 2 : NIH/3T3 (mouse embryo fibroblast cell line) whole cell lysate

Lane 3 : Mouse lung lysate

Lane 4 : Mouse spleen lysate

Lane 5 : Mouse testis lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Developed using the ECL technique.

Predicted band size: 18 kDa

Observed band size: 14,16 kDa

Exposure time: 3 seconds


Blocking/Dilution buffer: 5% NFDm/TBST.

Negative control: NIH/3T3 (PMID: 15210712).

Limited expression in mouse normal tissues has been documented in the literature (PMID: 9244355).

14 kDa band is a proteolytic p16INK4a lacking C-terminus (PMID: 18053084).

MEF cells are kindly provided by professor Pinlong Xu, Zhejiang University.



CDKN2A/p16INK4a was immunoprecipitated from 0.35 mg of MEF (mouse embryonic fibroblast cell line) whole cell lysate with ab211542 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab211542 at 1/5000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366), was used for detection at 1/10,000 dilution

Lane 1: MEF whole cell lysate 10 µg (Input).

Lane 2: ab211542 IP in MEF whole cell lysate.

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab211542 in MEF whole cell lysate.

Blocking/Dilution buffer: 5% NFDm/TBST.

Exposure time: 30 seconds.

MEF cells were kindly provided by professor Pinlong Xu, Zhejiang University.

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-CDKN2A/p16INK4a antibody [EPR20418] (ab211542)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors