

Product datasheet

Anti-Chlamydia trachomatis MOMP antibody
ab252752

Overview

Product name	Anti-Chlamydia trachomatis MOMP antibody
Description	Goat polyclonal to Chlamydia trachomatis MOMP
Host species	Goat
Specificity	Major Outer Membrane Protein (MOMP). Reacts with all serovars (A-K, L1-L3). Does not react with <i>C. psittacii</i> or <i>C. pneumoniae</i> in MIF. Negative against HEp-2 cells and egg yolk sac.
Tested applications	Suitable for: FM
Immunogen	Full length native protein (purified) corresponding to Chlamydia trachomatis MOMP. Strain L2.
General notes	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.1% Sodium azide Constituent: PBS
Purity	Ion Exchange Chromatography
Purification notes	> 95% pure. Sodium sulfate precipitation and ion-exchange chromatography.
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee

Our [Abpromise guarantee](#) covers the use of ab252752 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
FM		Use at an assay dependent concentration. Suitable for use in fluorescence microscopy. Also suitable for conjugation purposes. Each laboratory should determine an optimum working titer for use in its particular application.

Target

Relevance

Chlamydia is caused by the bacterium *Chlamydia trachomatis*. The intracytoplasmic inclusions caused by the bacterium are draped around the infected cell's nucleus. *Chlamydia trachomatis* is an intracellular organism that has a genome size of approximately 500-1000 kilobases and contains both RNA and DNA. The organism is also extremely temperature sensitive and must be refrigerated at 4°C as soon as a sample is obtained. Colonization of *Chlamydia* begins with attachment to sialic acid receptors on the eye, throat or genitalia. It persists at body sites that are inaccessible to phagocytes, T cells, and B cells. It also exists as 15 different serotypes. These serotypes cause four major diseases in humans: endemic trachoma (caused by serotypes A and C), sexually transmitted disease and inclusion conjunctivitis (caused by serotypes D and K), and lymphogranuloma venereum (caused by serotypes L1, L2, and L3). Studies reveal that *Chlamydia*, because of its cell wall, is able to inhibit phagolysosome fusion in phagocytes. The cell wall is proposed to be Gram negative in that it contains an outer lipopolysaccharide membrane, but it lacks peptidoglycan in its cell wall. This lack of peptidoglycan is shown by the inability to detect muramic acid and antibodies directed against it. It may, however, contain a carboxylated sugar other than muramic acid. The proposed structure consists of a major outer membrane protein cross linked with disulfide bonds. It also contains cysteine rich proteins (CRP) that may be the functional equivalent to peptidoglycan. This unique structure allows for intracellular division and extracellular survival (Hatch 1996). *Chlamydia* usually infects the cervix and fallopian tubes of women and the urethra of men. Chlamydial infections are believed to be one of the most common of all STDs. It is generally thought that in a population of 15 million, there are up to 300,000 cases of chlamydia each year. Thus, there are many undiagnosed cases of chlamydia in the community. It has been estimated that the true prevalence of chlamydia in the sexually active population may be in the order of 5% to 10%. *Chlamydia* is one of the leading causes of blindness in underdeveloped countries.

Cellular localization

Outer membrane; multi pass membrane protein.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors