

Product datasheet

Anti-Dnmt3a antibody [EPR18455] ab188470

KO VALIDATED Recombinant RabMAB

★★★★☆ 2 Abreviews 4 References 7 Images

Overview

| | |
|----------------------------|---|
| Product name | Anti-Dnmt3a antibody [EPR18455] |
| Description | Rabbit monoclonal [EPR18455] to Dnmt3a |
| Host species | Rabbit |
| Tested applications | Suitable for: Flow Cyt, WB, ICC/IF |
| Species reactivity | Reacts with: Mouse, Rat, Human |
| Immunogen | Synthetic peptide (the amino acid sequence is considered to be commercially sensitive) within Human Dnmt3a aa 600-700. The exact sequence is proprietary. Database link: Q9Y6K1 |
| Positive control | WB: HeLa, HEK-293 and C6 cell lysates; Rat brain and heart lysates. ICC/IF: HeLa cells. |
| General notes | Our RabMAB [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAB[®] patents . This product is a recombinant rabbit monoclonal antibody . |

Properties

| | |
|-----------------------------|---|
| Form | Liquid |
| Storage instructions | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. |
| Storage buffer | Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA |
| Purity | Protein A purified |
| Clonality | Monoclonal |
| Clone number | EPR18455 |
| Isotype | IgG |

Applications

Our [Abpromise guarantee](#) covers the use of **ab188470** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| Flow Cyt | | Use at an assay dependent concentration. |
| WB | ★★★★☆ | 1/2000. Detects a band of approximately 130 kDa (predicted molecular weight: 102 kDa). |
| ICC/IF | | 1/1000. |

Target

| | |
|---|--|
| Function | Required for genome wide de novo methylation and is essential for the establishment of DNA methylation patterns during development. DNA methylation is coordinated with methylation of histones. It modifies DNA in a non-processive manner and also methylates non-CpG sites. May preferentially methylate DNA linker between 2 nucleosomal cores and is inhibited by histone H1. Plays a role in paternal and maternal imprinting. Required for methylation of most imprinted loci in germ cells. Acts as a transcriptional corepressor for ZNF238. Can actively repress transcription through the recruitment of HDAC activity. |
| Tissue specificity | Highly expressed in fetal tissues, skeletal muscle, heart, peripheral blood mononuclear cells, kidney, and at lower levels in placenta, brain, liver, colon, spleen, small intestine and lung. |
| Sequence similarities | Belongs to the C5-methyltransferase family. Contains 1 ADD domain. Contains 1 GATA-type zinc finger. Contains 1 PHD-type zinc finger. Contains 1 PWWP domain. |
| Domain | The PWWP domain is essential for targeting to pericentric heterochromatin. |
| Post-translational modifications | Sumoylated; sumoylation disrupts the ability to interact with histone deacetylases (HDAC1 and HDAC2) and repress transcription. |
| Cellular localization | Nucleus. Cytoplasm. Accumulates in the major satellite repeats at pericentric heterochromatin. |

Images



Western blot - Anti-Dnmt3a antibody [EPR18455] (ab188470)

Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: Dnmt3a knockout HAP1 cell lysate (20 µg)

Lane 3: HeLa cell lysate (20 µg)

Lane 4: Mouse brain tissue lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab188470 observed at 125 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

ab188470 was shown to recognize Dnmt3a when Dnmt3a knockout samples were used, along with additional cross-reactive bands. Wild-type and Dnmt3a knockout samples were subjected to SDS-PAGE. ab188470 and [ab8245](#) (loading control to GAPDH) were diluted to 1/5000 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Dnmt3a antibody [EPR18455] (ab188470)

All lanes : Anti-Dnmt3a antibody [EPR18455] (ab188470) at 1/5000 dilution

Lane 1 : HeLa whole cell lysate

Lane 2 : Mouse skin tissue lysate

Lane 3 : Rat skin tissue lysate

Lane 4 : Mouse brain tissue lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

Performed under reducing conditions.


Predicted band size: 102 kDa

Observed band size: 125 kDa

[why is the actual band size different from the predicted?](#)


Lanes 1 - 4: Merged signal (red and green). Green - ab188470 observed at 125 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using Licor blocking buffer before being incubated with ab188470 and ab8245 (loading control) overnight at 4°C. Antibody binding was detected using Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) at a 1:10000 dilution for 1hr at room temperature and then imaged.



Flow Cytometry - Anti-Dnmt3a antibody [EPR18455] (ab188470)

Flow Cytometry analysis of HeLa (human cervix adenocarcinoma) cells labeling Dnmt3a with purified ab188470 at 1/80 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488)(1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



Western blot - Anti-Dnmt3a antibody [EPR18455] (ab188470)

Anti-Dnmt3a antibody [EPR18455] (ab188470) at 1/10000 dilution + HeLa (Human epithelial cells from cervix adenocarcinoma) cell lysate at 20 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 102 kDa


Observed band size: 130 kDa [why is the actual band size different from the predicted?](#)

Exposure time: 5 seconds

Blocking/Dilution buffer: 5% NFDm/TBST.

The observed MW is consistent with what has been described in

the literature (J Biol Chem. 2002. 277, 38746-38754. PMID: 2138111).



Western blot - Anti-Dnmt3a antibody [EPR18455] (ab188470)

Anti-Dnmt3a antibody [EPR18455] (ab188470) at 1/10000 dilution + HEK-293 (Human epithelial cells from embryonic kidney) cell lysate at 20 μ g

Secondary

Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution


Predicted band size: 102 kDa

Observed band size: 130 kDa [why is the actual band size different from the predicted?](#)

Exposure time: 15 seconds

Blocking/Dilution buffer: 5% NFDM/TBST.

The observed MW is consistent with what has been described in the literature (J Biol Chem. 2002. 277, 38746-38754. PMID: 2138111).



Western blot - Anti-Dnmt3a antibody [EPR18455] (ab188470)

All lanes : Anti-Dnmt3a antibody [EPR18455] (ab188470) at 1/2000 dilution

Lane 1 : Rat brain lysate

Lane 2 : Rat heart lysate

Lane 3 : C6 (Rat glial tumor cells) cell lysate

Lysates/proteins at 10 μ g per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 102 kDa

Observed band size: 130 kDa [why is the actual band size different from the predicted?](#)

Exposure time: 30 seconds

Blocking/Dilution buffer: 5% NFD/MTBST.



Immunocytochemistry/ Immunofluorescence - Anti-Dnmt3a antibody [EPR18455] (ab188470)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cells from cervix adenocarcinoma) cells labeling Dnmt3a with ab188470 at 1/1000 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green). Confocal image showing nuclear and weakly cytoplasmic staining on HeLa cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).

The negative controls are as follows:

-ve control 1: ab188470 at 1/1000 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.
-ve control 2: ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors