

Product datasheet

Anti-ERK1 + ERK2 antibody [EPR17526] ab184699

KO VALIDATED Recombinant RabMAB

★★★★★ [1 Abreviews](#) [121 References](#) [11 Images](#)

Overview

Product name	Anti-ERK1 + ERK2 antibody [EPR17526]
Description	Rabbit monoclonal [EPR17526] to ERK1 + ERK2
Host species	Rabbit
Tested applications	Suitable for: Flow Cyt (Intra), IP, ICC/IF, WB
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
Positive control	WB: Human ERK1 full length recombinant protein; Human ERK2 full length recombinant protein; A431, Jurkat, HeLa, HepG2, C6, RAW 264.7, PC-12 and NIH/3T3 whole cell lysates; Human fetal brain, fetal heart and fetal kidney lysates; Mouse brain, heart, kidney and spleen lysates; Rat brain, heart, kidney and spleen lysates. ICC/IF: HeLa cells. Flow Cyt (intra): A431 cells. IP: PC-12 whole cell extract.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAB[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAB[®] patents.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal

Clone number EPR17526

Isotype IgG

Applications

The Abpromise guarantee Our **Abpromise guarantee** covers the use of ab184699 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		1/440. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IP		1/70.
ICC/IF	★★★★★ (1)	1/250.
WB		1/10000. Detects a band of approximately 44, 42 kDa (predicted molecular weight: 43, 41 kDa).

Target

Function

Involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK1. Phosphorylates EIF4EBP1; required for initiation of translation. Phosphorylates microtubule-associated protein 2 (MAP2). Phosphorylates SPZ1 (By similarity). Phosphorylates heat shock factor protein 4 (HSF4) and ARHGEF2. Acts as a transcriptional repressor. Binds to a [GC]AAA[GC] consensus sequence. Repress the expression of interferon gamma-induced genes. Seems to bind to the promoter of CCL5, DMP1, IFIH1, IFITM1, IRF7, IRF9, LAMP3, OAS1, OAS2, OAS3 and STAT1. Transcriptional activity is independent of kinase activity.

Sequence similarities

Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily. Contains 1 protein kinase domain.

Domain

The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases.

Post-translational modifications

Dually phosphorylated on Thr-185 and Tyr-187, which activates the enzyme. Dephosphorylated by PTPRJ at Tyr-187.


Cellular localization

Nucleus.

Form

Mainly expressed in the cytoplasm and only localizes to the nucleus with treatment.

Images



All lanes : Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699) at 1/10000 dilution

Lane 1 : Mouse brain lysates

Lane 2 : Mouse heart lysates

Lane 3 : Mouse kidney lysates

Lane 4 : Mouse spleen lysates

Lane 5 : Rat brain lysates

Lane 6 : Rat heart lysates

Lane 7 : Rat kidney lysates

Lane 8 : Rat spleen lysates

Lane 9 : C6 (Rat glial tumor cells) whole cell lysates

Lane 10 : RAW 264.7 (Mouse macrophage cells transformed with Abelson murine leukemia virus) whole cell lysates

Lane 11 : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysates

Lane 12 : NIH/3T3 (Mouse embryo fibroblast cells) whole cell lysates

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 43, 41 kDa

Observed band size: 44, 42 kDa

44kDa band represents ERK1. 42kDa band represents ERK2.

Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)

All lanes : Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699) at 1/10000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : MAPK1 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 43, 41 kDa

Observed band size: 44 kDa

Lanes 1-2: Merged signal (red and green). Green - ab184699 observed at 44 kDa. Red - loading control **ab8245** observed at 37 kDa.


ab184699 Anti-ERK1 + ERK2 antibody [EPR17526] was shown to specifically react with ERK2 in wild-type HeLa cells. Loss of signal was observed when knockout cell line **ab265052** (knockout cell lysate **ab257525**) was used. Wild-type and ERK2 knockout samples were subjected to SDS-PAGE. ab184699 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 10000 Dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)

This image is courtesy of an Abreview submitted by Kirk Mcmanus.

Ab184699 staining ERK1 + ERK2 in HeLa cells by ICC/IF (Immunocytochemistry/Immunofluorescence). Cells were fixed with paraformaldehyde and permeabilized with 0.5% Triton X-100. Samples were incubated with primary antibody (1/500 in PBS) for 1 hour at 22°C. A Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed ([ab150081](#)) was used as the secondary antibody.



Western blot - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)

Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699) at 1/10000 dilution + Recombinant Human ERK1 protein ([ab43623](#)) ([ab43623](#)) at 0.01 µg

Secondary


Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 43, 41 kDa

Observed band size: 44 kDa

Recombinant full length ERK1 protein ([ab43623](#)) contains aa1-379.

Blocking/Dilution buffer: 5% NFDM/TBST.




Immunocytochemistry/ Immunofluorescence - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cells from cervix adenocarcinoma) cells labeling ERK1 + ERK2 with ab184699 at 1/250 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/500 dilution (green). Confocal image showing both nuclear and cytoplasmic staining on HeLa cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:

-ve control 1: ab184699 at 1/250 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.

-ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.



Western blot - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)

Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699) at 1/10000 dilution + Recombinant Human ERK2 protein (**ab43625**) at 0.01 µg

Secondary


Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 43, 41 kDa

Observed band size: 42 kDa

Recombinant full length ERK2 protein (**ab43625**) contains aa1-360.

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)

All lanes : Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699) at 1/50000 dilution

Lane 1 : A431 (Human epidermoid carcinoma) whole cell lysates

Lane 2 : Jurkat (Human T cell leukemia cells from peripheral blood) whole cell lysates

Lane 3 : HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysates

Lane 4 : HepG2 (Human liver hepatocellular carcinoma) whole cell lysates

Lane 5 : Human fetal brain lysates

Lane 6 : Human fetal heart lysates

Lane 7 : Human fetal kidney lysates

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 43, 41 kDa

Observed band size: 42,44 kDa

44kDa band represents ERK1. 42kDa band represents ERK2.

Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)

All lanes : Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699) at 1/10000 dilution

Lane 1 : Human fetal brain lysates

Lane 2 : Human fetal heart lysates

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 43, 41 kDa

Observed band size: 42,44 kDa


44kDa band represents ERK1. 42kDa band represents ERK2.

Blocking/Dilution buffer: 5% NFDm/TBST.



Intracellular flow cytometric analysis of A431 (Human epidermoid carcinoma) cells labeling ERK1 + ERK2 with ab184699 at 1/440 dilution (red) compared with a rabbit monoclonal IgG isotype control (black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody.

Flow Cytometry (Intracellular) - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)



ERK1 + ERK2 were immunoprecipitated from 1mg of PC-12 (Rat adrenal gland pheochromocytoma) whole cell extract with ab184699 at 1/70 dilution. Western blot was performed from the immunoprecipitate using ab184699 at 1/5000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500 dilution.

Lane 1: PC-12 whole cell extract. Lane 2: PBS instead of PC-12 whole cell extract.

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

44kDa band represents ERK1. 42kDa band represents ERK2.

Immunoprecipitation - Anti-ERK1 + ERK2 antibody [EPR17526] (ab184699)

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-ERK1 + ERK2 antibody [EPR17526]
(ab184699)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors