

Product datasheet

Anti-ERK5 antibody [EP791Y] - BSA and Azide free ab232538

KO VALIDATED Recombinant RabMAb[®]

5 Images

Overview

Product name	Anti-ERK5 antibody [EP791Y] - BSA and Azide free
Description	Rabbit monoclonal [EP791Y] to ERK5 - BSA and Azide free
Host species	Rabbit
Tested applications	Suitable for: WB, IP, ICC/IF, Flow Cyt
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide within Human ERK5 aa 800-900 (C terminal). The exact sequence is proprietary.
Positive control	WB: HAP1 and HeLa whole cell lysate.
General notes	Ab232538 is the carrier-free version of ab40809 . This format is designed for use in antibody labeling, including fluorochromes, metal isotopes, oligonucleotides, enzymes.

Our [carrier-free formats](#) are supplied in a buffer free of BSA, sodium azide and glycerol for higher conjugation efficiency.

Use our [conjugation kits](#) for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

ab232538 is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm.

Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	Constituent: PBS
Carrier free	Yes
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EP791Y
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab232538** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent concentration. Detects a band of approximately 115 kDa (predicted molecular weight: 88 kDa).
IP		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Target

Function	Plays a role in various cellular processes such as proliferation, differentiation and cell survival. The upstream activator of MAPK7 is the MAPK kinase MAP2K5. Upon activation, it translocates to the nucleus and phosphorylates various downstream targets including MEF2C. EGF activates MAPK7 through a Ras-independent and MAP2K5-dependent pathway. May have a role in muscle cell differentiation. May be important for endothelial function and maintenance of blood vessel integrity. MAP2K5 and MAPK7 interact specifically with one another and not with MEK1/ERK1 or MEK2/ERK2 pathways.
Tissue specificity	Expressed in many adult tissues. Abundant in heart, placenta, lung, kidney and skeletal muscle. Not detectable in liver.
Sequence similarities	Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily. Contains 1 protein kinase domain.
Domain	The second proline-rich region may interact with actin targeting the kinase to a specific location in the cell. The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases.


Post-translational modifications

Dually phosphorylated on Thr-219 and Tyr-221, which activates the enzyme (By similarity). Autophosphorylated in vitro on threonine and tyrosine residues when the C-terminal part of the kinase, which could have a regulatory role, is absent.

Cellular localization

Cytoplasm. Nucleus. Translocates to the nucleus upon activation.

Images



Western blot - Anti-ERK5 antibody [EP791Y] - BSA and Azide free (ab232538)

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)


Lane 2: MAPK7 (ERK5) knockout HAP1 whole cell lysate (20 µg)

Lane 3: HeLa whole cell lysate (20 µg)

Lanes 1 - 3: Merged signal (red and green). Green - [ab40809](#) observed at 88 kDa. Red - loading control, [ab9484](#), observed at 37 kDa.

[ab40809](#) was shown to specifically react with ERK5 in wild-type HAP1 cells as signal was lost in MAPK7 (ERK5) knockout cells. Wild-type and MAPK7 (ERK5) knockout samples were subjected to SDS-PAGE. [Ab40809](#) and [ab9484](#) (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab40809](#)).



Immunoprecipitation - Anti-ERK5 antibody [EP791Y] - BSA and Azide free (ab232538)

[ab40809](#) (purified) at 1:30 dilution (2ug) immunoprecipitating ERK5 in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10ug


Lane 2 (+): [ab40809](#) & HeLa whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of [ab40809](#) in HeLa whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDMM/TBST.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab40809](#)).



Flow Cytometry - Anti-ERK5 antibody [EP791Y] - BSA and Azide free (ab232538)

Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling ERK5 with purified [ab40809](#) at 1:50 dilution (10 ug/ml) (red). Cells were fixed with 80% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1:2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab40809](#)).



Immunocytochemistry/ Immunofluorescence - Anti-ERK5 antibody [EP791Y] - BSA and Azide free (ab232538)

Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling ERK5 with Purified [ab40809](#) at 1:100 (4.8 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab40809](#)).



Flow Cytometry - Anti-ERK5 antibody [EP791Y] - BSA and Azide free (ab232538)

Overlay histogram showing A549 cells stained with unpurified [ab40809](#) (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody ([ab40809](#), 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) ([ab150077](#)) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab40809](#)).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors