

Product datasheet

Anti-EYA4 antibody ab93865

[2 Images](#)

Overview

Product name	Anti-EYA4 antibody
Description	Rabbit polyclonal to EYA4
Host species	Rabbit
Tested applications	Suitable for: WB, IP
Species reactivity	Reacts with: Human Predicted to work with: Rabbit, Guinea pig, Chimpanzee, Rhesus monkey, Gorilla, Orangutan



Immunogen

Synthetic peptide from within residues: MEDSQLNEQ SVKKTCTESD VSQSQNSRSM EMQDLASPHT LVGGGDTPGS, corresponding to N terminal amino acids 1-50 of Human EYA4 (NP_004091.3)

[Run BLAST with ExPASy](#) [Run BLAST with NCBI](#)

Positive control

HeLa whole cell lysate

General notes

Reproducibility is key to advancing scientific discovery and accelerating scientists' next breakthrough.

Abcam is leading the way with our range of recombinant antibodies, knockout-validated antibodies and knockout cell lines, all of which support improved reproducibility.

We are also planning to innovate the way in which we present recommended applications and species on our product datasheets, so that only applications & species that have been tested in our own labs, our suppliers or by selected trusted collaborators are covered by our Abpromise™ guarantee.

In preparation for this, we have started to update the applications & species that this product is Abpromise guaranteed for.

We are also updating the applications & species that this product has been "predicted to work with," however this information is not covered by our Abpromise guarantee.

Applications & species from publications and Abreviews that have not been tested in our own labs or in those of our suppliers are not covered by the Abpromise guarantee.

Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, as well as customer reviews and Q&As.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	pH: 7 Preservative: 0.09% Sodium azide Constituent: Tris citrate/phosphate
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab93865** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
WB		
IP		

Application notes	IP: Use at 2-5µg/mg of lysate. WB: 1/1000 - 1/5000. Predicted molecular weight: 70 kDa. Not yet tested in other applications. Optimal dilutions/concentrations should be determined by the end user.
--------------------------	---

Target

Function	Tyrosine phosphatase that specifically dephosphorylates 'Tyr-142' of histone H2AX (H2AXY142ph). 'Tyr-142' phosphorylation of histone H2AX plays a central role in DNA repair and acts as a mark that distinguishes between apoptotic and repair responses to genotoxic stress. Promotes efficient DNA repair by dephosphorylating H2AX, promoting the recruitment of DNA repair complexes containing MDC1. Its function as histone phosphatase probably explains its role in transcription regulation during organogenesis. May be involved in development of the eye.
Tissue specificity	Highly expressed in heart and skeletal muscle.
Involvement in disease	Defects in EYA4 are the cause of deafness autosomal dominant type 10 (DFNA10) [MIM:601316]. DFNA10 is a form of sensorineural hearing loss. Sensorineural deafness results from damage to the neural receptors of the inner ear, the nerve pathways to the brain, or the area of the brain that receives sound information. Defects in EYA4 are the cause of cardiomyopathy dilated type 1J (CMD1J) [MIM:605362]. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death.
Sequence similarities	Belongs to the HAD-like hydrolase superfamily. EYA family.

Images



Western blot - Anti-EYA4 antibody (ab93865)

All lanes : Anti-EYA4 antibody (ab93865) at 0.4 $\mu\text{g/ml}$

Lane 1 : HeLa cell lysate at 50 μg


Lane 2 : HeLa cell lysate at 15 μg

Lane 3 : HeLa cell lysate at 5 μg

Developed using the ECL technique.

Predicted band size: 70 kDa

Exposure time: 3 minutes



Immunoprecipitation - Anti-EYA4 antibody (ab93865)

3 μg ab93865 were used to immunoprecipitate EYA4 from 1mg HeLa whole cell lysate. 20% IP was loaded and probed with 1 $\mu\text{g/ml}$ ab93865.

Detection: chemiluminescence with exposure time of 10 seconds.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors