

Product datasheet

Anti-FGF9/GAF antibody [EPR19937] ab206408

Recombinant RabMAb

4 Images

Overview

<b>Product name</b>	Anti-FGF9/GAF antibody [EPR19937]
<b>Description</b>	Rabbit monoclonal [EPR19937] to FGF9/GAF
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> WB
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	Recombinant fragment within Human FGF9/GAF aa 1 to the C-terminus. The exact sequence is proprietary. Mature protein without the proprotein from aa 1-3. Database link: <a href="#">P31371</a>
<b>Positive control</b>	WB: Human FGF9/GAF active protein; human fetal brain, fetal kidney and ovary cancer lysates; mouse kidney and brain lysates; rat brain lysate; C6, PC-12, NIH/3T3 and U-87 MG whole cell lysates.
<b>General notes</b>	Previously labelled as FGF9.  This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> For more information <a href="#">see here</a> .  Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a> .

Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	Preservative: 0.01% Sodium azide Constituents: 59% PBS, 0.05% BSA, 40% Glycerol
<b>Purity</b>	Protein A purified

<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EPR19937
<b>Isotype</b>	IgG

## Applications

---

Our [Abpromise guarantee](#) covers the use of **ab206408** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000. Detects a band of approximately 26 kDa (predicted molecular weight: 23 kDa).

---

## Target


---

<b>Function</b>	May have a role in glial cell growth and differentiation during development, gliosis during repair and regeneration of brain tissue after damage, differentiation and survival of neuronal cells, and growth stimulation of glial tumors.
<b>Tissue specificity</b>	Glial cells.
<b>Involvement in disease</b>	Defects in FGF9 are the cause of multiple synostoses syndrome type 3 (SYNS3) [MIM:612961]. Multiple synostoses syndrome is an autosomal dominant condition characterized by progressive joint fusions of the fingers, wrists, ankles and cervical spine, characteristic facies and progressive conductive deafness.
<b>Sequence similarities</b>	Belongs to the heparin-binding growth factors family.
<b>Post-translational modifications</b>	Three molecular species were found (30 kDa, 29 kDa and 25 kDa), cleaved at Leu-4, Val-13 and Ser-34 respectively. The smaller ones might be products of proteolytic digestion. Furthermore, there may be a functional signal sequence in the 30 kDa species which is uncleavable in the secretion step. N-glycosylated.
<b>Cellular localization</b>	Secreted.

---

## Images

---



Western blot - Anti-FGF9/GAF antibody [EPR19937] (ab206408)

**All lanes :** Anti-FGF9/GAF antibody [EPR19937] (ab206408) at 1/1000 dilution

**Lane 1 :** Human FGF9/GAF active protein

**Lane 2 :** Human FGF16 recombinant protein

**Lane 3 :** Human FGF20 recombinant protein

Lysates/proteins at 0.01 µg per lane.

**Secondary**

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution


**Predicted band size:** 23 kDa

**Observed band size:** 23 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.

Exposure times: Lane 1/2: 1 seconds; Lane 3: 15 seconds.

Human FGF9/GAF active protein contains aa1-208; FGF16 and FGF20 recombinant protein contain aa1-203 and aa4-208 respectively, both with His-Tag®.



Western blot - Anti-FGF9/GAF antibody [EPR19937] (ab206408)

**All lanes :** Anti-FGF9/GAF antibody [EPR19937] (ab206408) at 1/1000 dilution

**Lane 1 :** Human fetal brain lysate

**Lane 2 :** Human fetal kidney lysate

**Lane 3 :** Human ovary cancer lysate

Lysates/proteins at 10 µg per lane.

**Secondary**

**All lanes :** Goat Anti-Rabbit IgG Peroxidase Conjugate, specific to the non-reduced form of IgG at 1/100000 dilution

**Predicted band size:** 23 kDa


**Observed band size:** 26 kDa

why is the actual band size different from the predicted?

Blocking/Dilution buffer: 5% NFDM/TBST.

Exposure times: Lane 1/2: 15 seconds; Lane 3: 30 seconds.

The MW observed is consistent with the literature (PMID:16452189, PMID: 25925261).



**All lanes** : Anti-FGF9/GAF antibody [EPR19937] (ab206408) at 1/1000 dilution

**Lane 1** : Mouse kidney lysate

**Lane 2** : Mouse brain lysate

**Lane 3** : Rat brain lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes** : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution


**Predicted band size:** 23 kDa

**Observed band size:** 26 kDa [why is the actual band size different from the predicted?](#)

Blocking/Dilution buffer: 5% NFDm/TBST.

Exposure times: Lane 1/2: 15 seconds; Lane 3: 30 seconds.

The MW observed is consistent with the literature (PMID:16452189, PMID: 25925261).



**All lanes** : Anti-FGF9/GAF antibody [EPR19937] (ab206408) at 1/1000 dilution

**Lane 1** : C6 (Rat glioma cell line) whole cell lysate

**Lane 2** : PC-12 (Rat adrenal gland pheochromocytoma cell line) whole cell lysate

**Lane 3** : NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate

**Lane 4** : U-87 MG (Human glioblastoma-astrocytoma epithelial cell line) whole cell lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes** : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at

1/100000 dilution

**Predicted band size:** 23 kDa

**Observed band size:** 26 kDa [why is the actual band size different from the predicted?](#)

Blocking/Dilution buffer: 5% NFDm/TBST.

Exposure times: Lane 1-3: 15 seconds; Lane 4: 3 minutes.

The MW observed is consistent with the literature (PMID:16452189, PMID: 25925261).

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors