abcam

Product datasheet

Alexa Fluor® 488 Anti-Fibronectin antibody [F14] ab246735

Recombinant RabMAb

3 Images

Overview

Product name Alexa Fluor® 488 Anti-Fibronectin antibody [F14]

Description Alexa Fluor® 488 Rabbit monoclonal [F14] to Fibronectin

Host species Rabbit

Conjugation Alexa Fluor® 488, Ex: 495nm, Em: 519nm

Tested applications Suitable for: Flow Cyt, ICC/IF

Species reactivity Reacts with: Human

Predicted to work with: Mouse, Rat

Recombinant full length protein corresponding to Human Fibronectin aa 1-2400. **Immunogen**

Database link: P02751

Positive control ICC/IF: HepG2 cells. Flow Cytometry: HepG2 cells.

General notes This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Reproducibility is key to advancing scientific discovery and accelerating scientists' next breakthrough.

Abcam is leading the way with our range of recombinant antibodies, knockout-validated antibodies and knockout cell lines, all of which support improved reproducibility.

We are also planning to innovate the way in which we present recommended applications and species on our product datasheets, so that only applications & species that have been tested in our own labs, our suppliers or by selected trusted collaborators are covered by our Abpromise™ guarantee.

In preparation for this, we have started to update the applications & species that this product is Abpromise guaranteed for.

We are also updating the applications & species that this product has been "predicted to work with," however this information is not covered by our Abpromise guarantee.

Applications & species from publications and Abreviews that have not been tested in our own labs or in those of our suppliers are not covered by the Abpromise guarantee.

Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, as well as customer reviews and Q&As.

Properties

Form Liquid

Storage instructions Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle. Stable for 12 months at -20°C. Store In the Dark.

Storage buffer pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol, 1% BSA, PBS

Purity Protein A purified

Clonality Monoclonal

Clone number F14
Isotype IgG

Applications

Our Abpromise guarantee covers the use of ab246735 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application Abreviews	Notes
Flow Cyt	1/2500.

Application	Abreviews	Notes
ICC/IF		1/1000. This product gave a positive signal in HepG2 fixed with 4% formaldehyde (10 min) and 100% methanol (5 min).

Target

14.901	
Function	Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin. Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape. Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization. Participates in the regulation of type I collagen deposition by osteoblasts. Anastellin binds fibronectin and induces fibril formation. This fibronectin polymer, named superfibronectin, exhibits enhanced adhesive properties. Both anastellin and superfibronectin inhibit tumor growth, angiogenesis and metastasis. Anastellin activates p38 MAPK and inhibits lysophospholipid signaling.
Tissue specificity	Plasma FN (soluble dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine.
Involvement in disease	Glomerulopathy with fibronectin deposits 2
Sequence similarities	Contains 12 fibronectin type-I domains. Contains 2 fibronectin type-II domains. Contains 16 fibronectin type-III domains.
Developmental stage	Ugl-Y1, Ugl-Y2 and Ugl-Y3 are present in the urine from 0 to 17 years of age.
Post-translational modifications	Sulfated. It is not known whether both or only one of Thr-2064 and Thr-2065 are/is glycosylated. Forms covalent cross-links mediated by a transglutaminase, such as F13A or TGM2, between a glutamine and the epsilon-amino group of a lysine residue, forming homopolymers and heteropolymers (e.g. fibrinogen-fibronectin, collagen-fibronectin heteropolymers). Phosphorylated by FAM20C in the extracellular medium. Proteolytic processing produces the C-terminal NC1 peptide, anastellin.
Cellular localization	Secreted, extracellular space, extracellular matrix.

Images




Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-Fibronectin antibody [F14] (ab246735)

ab246735 staining Fibronectin in HepG2 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab246735 at 1/1000 dilution (shown in green) and ab195889, Mouse monoclonal to alpha Tubulin (Alexa Fluor®594), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HepG2 cells fixed with 100% methanol (5 min).




Flow Cytometry - Alexa Fluor® 488 Anti-Fibronectin antibody [F14] (ab246735)

Overlay histogram showing HepG2 cells stained with ab246735 (red line). The cells were fixed with 4 % formaldehyde (10 min) and then permeabilized with 0.1 % PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10 % normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab246735) (1x 10^6 in 100μ I at 0.2μ g/ml (1/2500)) for 30 min at 22° C.

Isotype control antibody (black line) was Rabbit IgG (monoclonal) Alexa Fluor 488[®] (ab199091) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter.

This antibody gave a positive signal inHepG2 cells fixed with 80 % methanol (5 min) / permeabilized with 0.1 % PBS-Triton X-100 for 15 min used under the same conditions.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.com/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors