

Product datasheet

Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] ab109364

Recombinant RabMAb

★★★★☆ 11 Abreviews 38 References 9 Images

Overview

Product name	Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805]
Description	Rabbit monoclonal [EPR4805] to GABARAP+GABARAPL1+GABARAPL2
Host species	Rabbit
Specificity	ab109364 will also recognize related targets GABARAPL1 and GABARAPL2: GABARAPL1: pBLAST 100% immunogen homology AND Abreview 38385 WB data GABARAPL2: pBLAST 93% homology (only 1aa gap of 14aa) AND Abreview 38385 WB data
Tested applications	Suitable for: Flow Cyt (Intra), WB, IHC-P, ICC/IF Unsuitable for: IP
Species reactivity	Reacts with: Mouse, Rat, Human Predicted to work with: Silk worm
Immunogen	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
Positive control	Fetal kidney lysate, fetal brain lysate, SH-SY5Y cell lysate, HepG2 whole cell lysate (ab7900), human brain tissue, HeLa cells.
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), PBS, 0.05% BSA
Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR4805
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab109364 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt (Intra)		1/20.
WB	★★★★★ (6)	1/1000 - 1/10000. Predicted molecular weight: 14 kDa.
IHC-P	★★★★★ (1)	1/500 - 1/1000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. Antigen retrieval is recommended. Ssee IHC antigen retrieval protocols . The mouse and rat recommendation is based on the WB results.
ICC/IF	★★★★★ (1)	1/100 - 1/500.

Application notes Is unsuitable for IP.

Target

Relevance


GABARAP: Ubiquitin-like modifier that plays a role in intracellular transport of GABA(A) receptors and its interaction with the cytoskeleton. Involved in apoptosis. Involved in autophagy. Whereas LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation.

GABARAPL1: Ubiquitin-like modifier that increases cell-surface expression of kappa-type opioid receptor through facilitating anterograde intracellular trafficking of the receptor. Involved in formation of autophagosomal vacuoles. Whereas LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation.

GABARAPL2: Ubiquitin-like modifier involved in intra-Golgi traffic. Modulates intra-Golgi transport through coupling between NSF activity and SNAREs activation. It first stimulates the ATPase activity of NSF which in turn stimulates the association with GOSR1 (By similarity). Involved in autophagy. Plays a role in mitophagy which contributes to regulate mitochondrial quantity and


quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production. Whereas LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation.

Images




Immunocytochemistry/ Immunofluorescence - Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] (ab109364)

Immunocytochemistry/Immunofluorescence analysis of HeLa (human cervix adenocarcinoma) labelling GABARAP+GABARAPL1+GABARAPL2 with purified ab109364 at 1/500. Cells were fixed with 4% PFA and permeabilized with 0.1% Triton X-100. An Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody (Ab150077). Nuclei counterstained with DAPI (blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] (ab109364)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human colon tissue sections labeling GABARAP+GABARAPL1+GABARAPL2 with Purified ab109364 at 1:500 dilution (0.29 µg/ml). Heat mediated antigen retrieval was performed using EDTA Buffer, pH 9.0. Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.



Flow Cytometry (Intracellular) - Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] (ab109364)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling GABARAP+GABARAPL1+GABARAPL2 with purified ab109364 at 1/20 dilution (10 ug/ml) (red). Cells were fixed with 4% Paraformaldehyde and permeabilized with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] (ab109364)

Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] (ab109364) at 1/5000 dilution (purified) + Rat kidney lysates at 20 µg


Secondary

Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 14 kDa

Observed band size: 17 kDa

Blocking and diluting buffer: 5% NFDM/TBST



Western blot - Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] (ab109364)

All lanes : Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] (ab109364) at 1/5000 dilution (purified)

Lane 1 : Mouse kidney lysates

Lane 2 : HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 14 kDa

Observed band size: 17 kDa

Blocking and diluting buffer: 5% NFDm/TBST




Flow Cytometry (Intracellular) - Anti-GABARAP+GABARAPL1+GABARAPL2 antibody [EPR4805] (ab109364)

Unpurified ab109364 staining

GABARAP+GABARAPL1+GABARAPL2 in the human cell line HeLa (human cervix adenocarcinoma) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde, permeabilized with 90% methanol and the sample was incubated with the primary antibody at a dilution of 1/20. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.

Isotype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)



Western blot - Anti-
GABARAP+GABARAPL1+GABARAPL2 antibody
[EPR4805] (ab109364)

All lanes : Anti-GABARAP+GABARAPL1+GABARAPL2 antibody
[EPR4805] (ab109364) at 1/1000 dilution (unpurified)

Lane 1 : Fetal kidney lysate


Lane 2 : SH-SY5Y cell lysate

Lane 3 : HepG2 cell lysate

Lane 4 : Fetal brain lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 14 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-
GABARAP+GABARAPL1+GABARAPL2 antibody
[EPR4805] (ab109364)

Unpurified ab109364, at 1/500, staining

GABARAP+GABARAPL1+GABARAPL2 in Human brain tissue by
immunohistochemistry.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-GABARAP+GABARAPL1+GABARAPL2
antibody [EPR4805] (ab109364)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors