


Product datasheet

Anti-GALNT12 antibody ab101358

1 Image

Overview

| | |
|----------------------------|---|
| Product name | Anti-GALNT12 antibody |
| Description | Rabbit polyclonal to GALNT12 |
| Host species | Rabbit |
| Tested applications | Suitable for: WB |
| Species reactivity | Reacts with: Human Predicted to work with: Mouse, Cow  |
| Immunogen | Recombinant fragment, corresponding to a region within internal sequence amino acids 34-173 of Human GALNT12 (AAH13945; UniProt ID: Q8IXK2 isoform 2). |
| Positive control | A549 whole cell lysate |
| General notes | <p>Reproducibility is key to advancing scientific discovery and accelerating scientists' next breakthrough.</p> <p>Abcam is leading the way with our range of recombinant antibodies, knockout-validated antibodies and knockout cell lines, all of which support improved reproducibility.</p> <p>We are also planning to innovate the way in which we present recommended applications and species on our product datasheets, so that only applications & species that have been tested in our own labs, our suppliers or by selected trusted collaborators are covered by our Abpromise™ guarantee.</p> <p>In preparation for this, we have started to update the applications & species that this product is Abpromise guaranteed for.</p> <p>We are also updating the applications & species that this product has been “predicted to work with,” however this information is not covered by our Abpromise guarantee.</p> <p>Applications & species from publications and Abreviews that have not been tested in our own labs or in those of our suppliers are not covered by the Abpromise guarantee.</p> <p>Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, as well as customer reviews and Q&As.</p> |

Properties

| | |
|-------------|--------|
| Form | Liquid |
|-------------|--------|

| | |
|-----------------------------|---|
| Storage instructions | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Storage buffer | pH: 7.00 Preservative: 0.01% Thimerosal (merthiolate) Constituents: 1.21% Tris, 0.75% Glycine, 20% Glycerol (glycerin, glycerine) |
| Purity | Immunogen affinity purified |
| Clonality | Polyclonal |
| Isotype | IgG |

Applications

Our [Abpromise guarantee](#) covers the use of **ab101358** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| Application | Abreviews | Notes |
|-------------|-----------|--|
| WB | | 1/500 - 1/3000. Predicted molecular weight: for isoform 2, 32 kDa. |

Target

| | |
|-------------------------------|---|
| Function | Catalyzes the initial reaction in O-linked oligosaccharide biosynthesis, the transfer of an N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor. Has activity toward non-glycosylated peptides such as Muc5AC, Muc1a and EA2, and no detectable activity with Muc2 and Muc7. Displays enzymatic activity toward the Gal-NAc-Muc5AC glycopeptide, but no detectable activity to mono-GalNAc-glycosylated Muc1a, Muc2, Muc7 and EA2. May play an important role in the initial step of mucin-type oligosaccharide biosynthesis in digestive organs. |
| Tissue specificity | Widely expressed at different levels of expression. Highly expressed in digestive organs such as small intestine, stomach, pancreas and colon. Expressed at intermediate level in testis, thyroid gland and spleen. Weakly expressed in whole brain, cerebral cortex, cerebellum, fetal brain, bone marrow, thymus, leukocytes, heart, skeletal muscle, liver, lung, esophagus, kidney, adrenal gland, mammary gland, uterus, placenta, ovary and prostate. |
| Pathway | Protein modification; protein glycosylation. |
| Involvement in disease | Defects in GALNT12 are a cause of susceptibility to colorectal cancer type 1 (CRCS1) [MIM:608812]. Colorectal cancer is a malignancy originating either in the colon or rectum or both. |
| Sequence similarities | Belongs to the glycosyltransferase 2 family. GalNAc-T subfamily. Contains 1 ricin B-type lectin domain. |
| Domain | There are two conserved domains in the glycosyltransferase region: the N-terminal domain (domain A, also called GT1 motif), which is probably involved in manganese coordination and substrate binding and the C-terminal domain (domain B, also called Gal/GalNAc-T motif), which is probably involved in catalytic reaction and UDP-Gal binding. The ricin B-type lectin domain binds to GalNAc and contributes to the glycopeptide specificity. |
| Cellular localization | Golgi apparatus membrane. |

Images



Anti-GALNT12 antibody (ab101358) at 1/1000 dilution + A549 whole cell lysate at 30 μ g

Predicted band size: for isoform 2, 32 kDa

12% SDS-PAGE

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors