

Product datasheet

Anti-GFP antibody [EPR14104] ab183734

Recombinant **RabMAb**

★★★★★ 22 Abreviews 9 References 9 Images

Overview

Product name	Anti-GFP antibody [EPR14104]
Description	Rabbit monoclonal [EPR14104] to GFP
Host species	Rabbit
Tested applications	Suitable for: WB, IHC-P, ICC/IF, Flow Cyt Unsuitable for: IP
Species reactivity	Reacts with: Species independent
Immunogen	Full length protein corresponding to <i>Aequorea victoria</i> GFP aa 1 to the C-terminus. Database link: P42212 Run BLAST with Run BLAST with
Positive control	WB: GFP transfected 293 cell lysate. IHC-P: GFP transgenic mouse colon tissue and GFP transgenic mouse liver tissue. ICC/IF: GFP transfected 293 cells. ICC-IF: GFP transfected NIH3T3 cells.
General notes	<p>On the basis of low sequence homology, ab183734 is predicted to show no or limited cross-reactivity to RFP, BFP, and CFP.</p> <p>For the best results in IHC, please use ab183734 on Formalin/PFA-fixed paraffin-embedded sections with heat mediated antigen retrieval.</p> <p>A trial size is available to purchase for this antibody.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p> <p>This product is a recombinant rabbit monoclonal antibody.</p>


Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

Dissociation constant (K_D)

K_D = 1.11 x 10⁻¹¹ M



Storage buffer

Preservative: 0.01% Sodium azide
Constituents: 59% PBS, 40% Glycerol, 0.05% BSA

Purity

Protein A purified

Clonality

Monoclonal

Clone number

EPR14104

Isotype

IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab183734** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★★	1/10000 - 1/50000. Detects a band of approximately 27 kDa (predicted molecular weight: 27 kDa).
IHC-P	★★★★★	1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See IHC antigen retrieval protocols .
ICC/IF	★★★★★	1/500.
Flow Cyt		Use at an assay dependent concentration.

Application notes

Is unsuitable for IP.

Target

Relevance

Function: Energy-transfer acceptor. Its role is to transduce the blue chemiluminescence of the protein aequorin into green fluorescent light by energy transfer. Fluoresces in vivo upon receiving energy from the Ca²⁺-activated photoprotein aequorin.

Subunit structure: Monomer.

Tissue specificity: Photocytes.

Post-translational modification: Contains a chromophore consisting of modified amino acid residues. The chromophore is formed by autocatalytic backbone condensation between Ser-65 and Gly-67, and oxidation of Tyr-66 to didehydrotyrosine. Maturation of the chromophore requires nothing other than molecular oxygen.


Biotechnological use: Green fluorescent protein has been engineered to produce a vast


number of variously colored mutants, fusion proteins, and biosensors. Fluorescent proteins and its mutated allelic forms, blue, cyan and yellow have become a useful and ubiquitous tool for making chimeric proteins, where they function as a fluorescent protein tag. Typically they tolerate N- and C-terminal fusion to a broad variety of proteins. They have been expressed in most known cell types and are used as a noninvasive fluorescent marker in living cells and organisms. They enable a wide range of applications where they have functioned as a cell lineage tracer, reporter of gene expression, or as a measure of protein-protein interactions. Can also be used as a molecular thermometer, allowing accurate temperature measurements in fluids. The measurement process relies on the detection of the blinking of GFP using fluorescence correlation spectroscopy.

Sequence similarities: Belongs to the GFP family.

Biophysicochemical properties: Absorption: Abs(max)=395 nm
Exhibits a smaller absorbance peak at 470 nm. The fluorescence emission spectrum peaks at 509 nm with a shoulder at 540 nm.

Images






Immunocytochemistry/ Immunofluorescence - Anti-GFP antibody [EPR14104] (ab183734)

ab183734 staining GFP in GFP-transfected NIH3T3 cells. The cells were fixed with 4% formaldehyde (10min) and then blocked in 1% BSA / 0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated with ab183734 at 1/500 dilution overnight at +4°C followed by incubation with ab150081, Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488), for 1 hour, at 1µg/ml.

Under identical experimental conditions, when compared to the basal level of GFP expression in transfected NIH3T3 cells, the cells upon which ab183734 was applied gave a stronger signal in the 488 channel, indicating that ab183734 is binding to GFP and therefore eliciting signal amplification.

ab183734 was also applied to non-GFP-transfected NIH3T3 cells, which produced no positive staining, indicating specificity for GFP. Nuclear DNA was labelled with 1.43µM DAPI (blue).



Western blot - Anti-GFP antibody [EPR14104] (ab183734)

All lanes : Anti-GFP antibody [EPR14104] (ab183734) at 1/10000 dilution (purified)

Lane 1 : GFP transfected 293 cell lysate

Lane 2 : Non-transfected 293 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 27 kDa

Observed band size: 25 kDa


[why is the actual band size different from the predicted?](#)

Blocking and dilution buffer: 5% NFDM /TBST.




Flow Cytometry - Anti-GFP antibody [EPR14104] (ab183734)

Flow Cytometry analysis of 293T (human embryonic kidney) transfected with human TNFRSF9 cells labeling GFP with purified ab183734 at 1/50 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 647)(1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) was used as the unlabeled control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFP antibody [EPR14104] (ab183734)

Immunohistochemical analysis of paraffin-embedded GFP transgenic mouse colon tissue (left) and normal mouse colon tissue (right) labeling GFP with unpurified ab183734 at 1/250 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counterstained with Hematoxylin.



Immunocytochemistry/ Immunofluorescence - Anti-GFP antibody [EPR14104] (ab183734)

Immunofluorescent analysis of 4% paraformaldehyde-fixed GFP transfected 293 cells labeling GFP with unpurified ab183734 at 1/500 dilution, followed by Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody at 1/200 dilution (green). Counterstained with Dapi (blue)



Western blot - Anti-GFP antibody [EPR14104] (ab183734)

All lanes : Anti-GFP antibody [EPR14104] (ab183734) at 1/10000 dilution (unpurified)

Lane 1 : GFP transfected 293 cell lysate

Lane 2 : Non-transfected 293 cell lysate


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution


Predicted band size: 27 kDa

Observed band size: 27 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFP antibody [EPR14104] (ab183734)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse pancreas tissue (negative control) labelling GFP with purified ab183734 at a dilution of 1/100. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFP antibody [EPR14104] (ab183734)

Immunohistochemical analysis of paraffin-embedded GFP transgenic mouse liver tissue (left) and normal mouse liver tissue (right) labelling GFP with unpurified ab183734 at 1/250 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counterstained with Hematoxylin.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors