

Product datasheet

Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)] ab188317

Recombinant RabMAb

★★★★★ 3 Abreviews 3 References 4 Images

Overview

Product name	Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)]
Description	Rabbit monoclonal [EPR930(2)] to Glucose Transporter GLUT4 (phospho S488)
Host species	Rabbit
Tested applications	Suitable for: WB, Dot blot
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide (the amino acid sequence is considered to be commercially sensitive) within Human Glucose Transporter GLUT4 aa 450 to the C-terminus. The exact sequence is proprietary. Database link: P14672
Positive control	Human fetal heart, mouse heart, rat heart
General notes	

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
Purity	Protein A purified

Clonality	Monoclonal
Clone number	EPR930(2)
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab188317** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★★	1/1000 - 1/10000. Detects a band of approximately 45 kDa (predicted molecular weight: 55 kDa).
Dot blot		Use at an assay dependent concentration.

Target

Function	Insulin-regulated facilitative glucose transporter.
Tissue specificity	Skeletal and cardiac muscles; brown and white fat.
Involvement in disease	Diabetes mellitus, non-insulin-dependent
Sequence similarities	Belongs to the major facilitator superfamily. Sugar transporter (TC 2.A.1.1) family. Glucose transporter subfamily.
Post-translational modifications	Sumoylated.
Cellular localization	Cell membrane. Endomembrane system. Cytoplasm, perinuclear region. Localizes primarily to the perinuclear region, undergoing continued recycling to the plasma membrane where it is rapidly reinternalized. The dileucine internalization motif is critical for intracellular sequestration.

Images



Western blot - Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)] (ab188317)

All lanes : Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)] (ab188317) at 1/1000 dilution

Lane 1 : Human fetal heart whole cell lysates

Lane 2 : Human fetal heart whole cell lysates with Phosphatase

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, Peroxidase conjugated (DC03L) at 1/20000 dilution


Predicted band size: 55 kDa

Observed band size: 45 kDa

why is the actual band size different from the predicted?

Exposure time: 1 second


Blocking/Diluting buffer and concentration 5% NFDm/TBST



Dot Blot - Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)] (ab188317)

Dot Blot analysis of Lane 1: Glucose Transporter GLUT4 (pS488) phospho peptide and Lane 2: Glucose Transporter GLUT4 non-phospho peptide labeling Glucose Transporter GLUT4 (phospho S488) with ab188317 at 1/1000 dilution.

5% NFDm/TBST was used as the diluting and blocking buffer. ab97051 Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated was used as the secondary antibody at 1/100000 dilution. Exposure time: 3 minutes.



Western blot - Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)] (ab188317)

All lanes : Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)] (ab188317) at 1/5000 dilution

Lane 1 : Mouse heart

Lane 2 : Rat heart


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 55 kDa

Observed band size: 45 kDa why is the actual band size different from the predicted?



Western blot - Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)] (ab188317)

All lanes : Anti-Glucose Transporter GLUT4 (phospho S488) antibody [EPR930(2)] (ab188317) at 1/20000 dilution

Lane 1 : Untreated human fetal heart

Lane 2 : Lambda phosphorylase treated human fetal heart

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 55 kDa

Observed band size: 45 kDa [why is the actual band size different from the predicted?](#)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors