

Product datasheet

Anti-GNE antibody ab244403

3 Images

Overview

Product name	Anti-GNE antibody
Description	Rabbit polyclonal to GNE
Host species	Rabbit
Tested applications	Suitable for: ICC/IF, WB, IHC-P
Species reactivity	Reacts with: Human Predicted to work with: Mouse, Rat, Chinese hamster 
Immunogen	Recombinant fragment corresponding to Human GNE aa 282-422. Sequence: VPFDQFIQLVAHAGCMIGNSSCGVREVGAFGTPVINLGTR QIGRETGENV LHVRDADTQDKILQALHLQFGKQYPCSKIYGDGNAVPRILK FLKSIDLQE PLQKKFCFPPVKENISQDIDHILETLSALAVDLGGTNLRVA Database link: Q9Y223  Run BLAST with  Run BLAST with
Positive control	ICC/IF: A431 cells. WB: A549 cell lysate. IHC-P: Human liver tissue.
General notes	This product was previously labelled as GLCNE

Reproducibility is key to advancing scientific discovery and accelerating scientists' next breakthrough.

Abcam is leading the way with our range of recombinant antibodies, knockout-validated antibodies and knockout cell lines, all of which support improved reproducibility.

We are also planning to innovate the way in which we present recommended applications and species on our product datasheets, so that only applications & species that have been tested in our own labs, our suppliers or by selected trusted collaborators are covered by our Abpromise™ guarantee.

In preparation for this, we have started to update the applications & species that this product is Abpromise guaranteed for.

We are also updating the applications & species that this product has been "predicted to work with," however this information is not covered by our Abpromise guarantee.

Applications & species from publications and Abreviews that have not been tested in our own

labs or in those of our suppliers are not covered by the Abpromise guarantee.

Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, as well as customer reviews and Q&As.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage buffer	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine)
Purity	Immunogen affinity purified
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab244403** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		Use a concentration of 0.25 - 2 µg/ml. Fixation/Permeabilization: PFA/Triton X-100.
WB		Use a concentration of 0.04 - 0.4 µg/ml.
IHC-P		1/50 - 1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Target

Function	Regulates and initiates biosynthesis of N-acetylneuraminic acid (NeuAc), a precursor of sialic acids. Plays an essential role in early development (By similarity). Required for normal sialylation in hematopoietic cells. Sialylation is implicated in cell adhesion, signal transduction, tumorigenicity and metastatic behavior of malignant cells.
Tissue specificity	Highest expression in liver and placenta. Also found in heart, brain, lung, kidney, skeletal muscle and pancreas. Isoform 1 is expressed in heart, brain, kidney, liver, placenta, lung, spleen, pancreas, skeletal muscle and colon. Isoform 2 is expressed mainly in placenta, but also in brain, kidney, liver, lung, pancreas and colon. Isoform 3 is expressed at low level in kidney, liver, placenta and colon.
Pathway	Amino-sugar metabolism; N-acetylneuraminate biosynthesis.

Involvement in disease

Defects in GNE are a cause of sialuria (SIALURIA) [MIM:269921]; also known as sialuria French type. In sialuria, free sialic acid accumulates in the cytoplasm and gram quantities of neuraminic acid are secreted in the urine. The metabolic defect involves lack of feedback inhibition of UDP-GlcNAc 2-epimerase by CMP-Neu5Ac, resulting in constitutive overproduction of free Neu5Ac. Clinical features include variable degrees of developmental delay, coarse facial features and hepatomegaly. Sialuria inheritance is autosomal dominant.

Defects in GNE are the cause of inclusion body myopathy type 2 (IBM2) [MIM:600737]. Hereditary inclusion body myopathies are a group of neuromuscular disorders characterized by adult onset, slowly progressive distal and proximal weakness and a typical muscle pathology including rimmed vacuoles and filamentous inclusions. IBM2 is an autosomal recessive disorder affecting mainly leg muscles, but with an unusual distribution that spares the quadriceps as also observed in Nonaka myopathy.

Defects in GNE are the cause of Nonaka myopathy (NM) [MIM:605820]; also known as distal myopathy with rimmed vacuoles (DMRV). NM is an autosomal recessive muscular disorder, allelic to inclusion body myopathy 2. It is characterized by weakness of the anterior compartment of the lower limbs with onset in early adulthood, and sparing of the quadriceps muscles. As the inclusion body myopathy, NM is histologically characterized by the presence of numerous rimmed vacuoles without inflammatory changes in muscle specimens.

Sequence similarities

In the N-terminal section; belongs to the UDP-N-acetylglucosamine 2-epimerase family.

In the C-terminal section; belongs to the ROK (NagC/XylR) family.


Post-translational modifications

Phosphorylated by PKC.

Cellular localization


Cytoplasm.

Images




PFA-fixed, Triton X-100 permeabilized A431 (human epidermoid carcinoma cell line) cells stained for GNE (green) using ab244403 at 4 μ g/ml in ICC/IF.

Immunocytochemistry/ Immunofluorescence - Anti-GNE antibody (ab244403)



Paraffin-embedded human liver tissue stained for GNE using ab244403 at 1/50 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GNE antibody (ab244403)



Anti-GNE antibody (ab244403) at 0.4 µg/ml + A549 (Human lung carcinoma cell line) cell lysate

Western blot - Anti-GNE antibody (ab244403)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors