

Product datasheet

Goat F(ab')₂ Anti-Mouse IgG - (Fab)'₂ (PE), pre-adsorbed ab5889

2 References

Overview

Product name	Goat F(ab') ₂ Anti-Mouse IgG - (Fab)' ₂ (PE), pre-adsorbed
Host species	Goat
Target species	Mouse
Tested applications	Suitable for: ICC/IF, Immunomicroscopy, Flow Cyt
Minimal cross-reactivity	Cow, Horse, Human, Rabbit, Rat, Sheep more details
Immunogen	Mouse IgG F(ab) ₂ fragment
Conjugation	PE. Ex: 488nm, Em: 575nm

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C.
Storage buffer	Preservative: 0.01% Sodium azide Constituents: 0.878% Sodium chloride, 1% BSA, 0.424% Potassium phosphate BSA is IgG and protease free
Purification notes	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation.
Conjugation notes	R-Phycoerythrin (R-PE) (Molecular Weight 240,000 daltons) from seaweed Absorption Wavelength: 490 nm, 545 nm and 565 nm Emission Wavelength: 580 nm OD _{566nm} /OD _{280nm} : 2.6 Phycobiliprotein Concentration: 0.5 mg/ml (determined by absorbance = 82.0 at 565 nm for a 1% solution for only those R-PE molecules to which at least one molecule of active antibody is bound).
Clonality	Polyclonal
Isotype	IgG

Applications

The Abpromise guarantee

Our [Abpromise guarantee](#) covers the use of ab5889 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
ICC/IF		1/100 - 1/250.
Immunomicroscopy		Use at an assay dependent dilution. Suitable for other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.
Flow Cyt		1/100 - 1/250. The maximum amount of reagent required to stain 1×10^6 cells in flow cytometry is approximately 1.0 μg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors