

## Product datasheet

# Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) ab150116

★★★★★ 3 Abreviews 195 References 5 Images

### Overview

<b>Product name</b>	Goat Anti-Mouse IgG H&L (Alexa Fluor® 594)
<b>Description</b>	Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594)
<b>Host species</b>	Goat
<b>Target species</b>	Mouse
<b>Tested applications</b>	<b>Suitable for:</b> IHC-Fr, ICC/IF, ELISA, IHC-P, Flow Cyt
<b>Immunogen</b>	The details of the immunogen for this antibody are not available.
<b>Conjugation</b>	Alexa Fluor® 594. Ex: 590nm, Em: 617nm

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Stable for 12 months at -20°C. Store In the Dark.
<b>Storage buffer</b>	Preservative: 0.02% Sodium azide Constituents: 23% Glycerol (glycerin, glycerine), PBS, 1% BSA
<b>Purity</b>	Immunogen affinity purified
<b>Purification notes</b>	The antibody was isolated by affinity chromatography using antigen coupled to agarose beads.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>General notes</b>	Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

## Applications


### The Abpromise guarantee

Our [Abpromise guarantee](#) covers the use of ab150116 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


Application	Abreviews	Notes
IHC-Fr	★★★★★ (1)	Use at an assay dependent concentration.
ICC/IF	★★★★★ (2)	1/200 - 1/1000.
ELISA		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.
Flow Cyt		1/2000 - 1/4000. <a href="#">ab178000</a> - Mouse monoclonal IgG1 (Alexa Fluor® 594), is suitable for use as an isotype control to complement this secondary antibody.

## Images



ICC/IF image of [ab7291](#) stained HeLa cells. The cells were 100% methanol fixed (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then incubated in 1% BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to block non-specific protein-protein interactions. The cells were then incubated with the antibody ([ab7291](#), 1µg/ml) overnight at +4°C. The secondary antibody (orange) was ab150116 Alexa Fluor® 594 goat anti-mouse IgG (H+L) used at 2µg/ml for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

The negative control (inset) is a secondary-only assay to demonstrate low non-specific binding of the secondary antibody.




Flow Cytometry - Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) (ab150116)

Overlay histogram showing Jurkat cells stained with [ab8090](#) (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody ([ab8090](#), 0.01µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody Goat anti-mouse IgG H&L (Alexa Fluor® 594) ([ab150116](#)) was used at 1/4000 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2a [ICIGG2A] ([ab91361](#), 0.01µg/1x10<sup>6</sup> cells) used under the same conditions.

Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 561nm laser and 610/20 bandpass filter.




ELISA - Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) (ab150116)

Cross-reactivity of the polyclonal secondary antibody [ab182017](#) was tested using a sandwich ELISA approach. The wells were coated with the indicated IgG standards at 1 µg/ml (50µl/well) and incubated overnight at 4°C, followed by a 5% BSA blocking step for 2h at RT. [ab182017](#) was then added starting at 1 µg/ml and gradually diluted 1/4 (50 µl/well), followed by incubation for 2h. For the detection Donkey anti-Goat IgG H&L (HRP) ([ab6885](#)) was used at 1/10,000 dilution (50 µl/well), followed by incubation for 1h at RT.

**For the batch tested, [ab182017](#) showed a cross-reactivity below 2% towards Chicken IgY, 6% towards Human IgG, 7% towards Rabbit IgG and 47% towards Rat IgG.**


This data was developed using the unconjugated antibody ([ab182017](#)).



ELISA - Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) (ab150116)

Cross-reactivity of Goat anti-Mouse IgG H&L ([ab182017](#)) and Goat anti-Mouse IgG H&L obtained from two different vendors was tested using a sandwich ELISA approach. The wells were coated with the indicated IgG standards (Rabbit, Human, Mouse and Rat) at 1 µg/ml (50 µl/well) and incubated overnight at 4°C, followed by a 5% BSA blocking step for 2h at RT. Secondary antibodies were then added starting at 1 µg/ml and gradually diluted 1/4 (50 µl/well), followed by incubation for 2h. For the detection Donkey anti-Goat IgG H&L (HRP) ([ab6885](#)) was used at 1/10,000 dilution (50 µl/well), followed by incubation for 1h at RT. This data is from a representative dilution.

This data was developed using the unconjugated antibody ([ab182017](#)).



Alexa Fluor® - Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) (ab150116)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors