

## Product datasheet

# Anti-Heparan Sulfate Proteoglycan 2/Perlecan antibody [A74] ab23418

★★★★☆ 6 Abreviews 12 References

### Overview

<b>Product name</b>	Anti-Heparan Sulfate Proteoglycan 2/Perlecan antibody [A74]
<b>Description</b>	Mouse monoclonal [A74] to Heparan Sulfate Proteoglycan 2/Perlecan
<b>Host species</b>	Mouse
<b>Specificity</b>	ab23418 is highly specific for perlecan. There is no evidence for cross-reactivity with other connective tissue proteins (vitronectin, fibronectin, elastin, collagen, laminin).
<b>Species reactivity</b>	<b>Reacts with:</b> Cow, Human
<b>Immunogen</b>	Tissue, cells or virus corresponding to Cow Heparan Sulfate Proteoglycan 2/Perlecan.
<b>Epitope</b>	Epitope is located in domain V of Perlecan.
<b>General notes</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	<p>pH: 7.40</p> <p>Preservative: 0.097% Sodium azide</p> <p>Constituents: 0.0268% PBS, 2.9% Sodium chloride</p>
<b>Purity</b>	Protein G purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	A74
<b>Myeloma</b>	Sp2/0
<b>Isotype</b>	IgG1

## Target

---

<b>Function</b>	<p>Integral component of basement membranes. Component of the glomerular basement membrane (GBM), responsible for the fixed negative electrostatic membrane charge, and which provides a barrier which is both size- and charge-selective. It serves as an attachment substrate for cells. Plays essential roles in vascularization. Critical for normal heart development and for regulating the vascular response to injury. Also required for avascular cartilage development.</p> <p>Endorepellin is an anti-angiogenic and anti-tumor peptide that inhibits endothelial cell migration, collagen-induced endothelial tube morphogenesis and blood vessel growth in the chorioallantoic membrane. Blocks endothelial cell adhesion to fibronectin and type I collagen. Anti-tumor agent in neovascularization. Interaction with its ligand, integrin alpha2/beta1, is required for the anti-angiogenic properties. Evokes a reduction in phosphorylation of receptor tyrosine kinases via alpha2/beta1 integrin-mediated activation of the tyrosine phosphatase, PTPN6.</p> <p>The LG3 peptide has anti-angiogenic properties that require binding of calcium ions for full activity.</p>
<b>Tissue specificity</b>	<p>Found in the basement membranes.</p>
<b>Involvement in disease</b>	<p>Defects in HSPG2 are the cause of Schwartz-Jampel syndrome (SJS1) [MIM:255800]; a rare autosomal recessive disorder characterized by permanent myotonia (prolonged failure of muscle relaxation) and skeletal dysplasia, resulting in reduced stature, kyphoscoliosis, bowing of the diaphyses and irregular epiphyses.</p> <p>Defects in HSPG2 are the cause of dyssegmental dysplasia Silverman-Handmaker type (DDSH) [MIM:224410]. The dyssegmental dysplasias are rare, autosomal recessive skeletal dysplasias with anisospondyly and micromelia. There are two recognized types: the severe, lethal DDSH and the milder Rolland-Desbuquois form. Individuals with DDSH also have a flat face, micrognathia, cleft palate and reduced joint mobility, and frequently have an encephalocoele. The endochondral growth plate is short, the calcospherites (which are spherical calcium-phosphorus crystals produced by hypertrophic chondrocytes) are unfused, and there is mucoid degeneration of the resting cartilage.</p>
<b>Sequence similarities</b>	<p>Contains 4 EGF-like domains.</p> <p>Contains 22 Ig-like C2-type (immunoglobulin-like) domains.</p> <p>Contains 11 laminin EGF-like domains.</p> <p>Contains 3 laminin G-like domains.</p> <p>Contains 3 laminin IV type A domains.</p> <p>Contains 4 LDL-receptor class A domains.</p> <p>Contains 1 SEA domain.</p>
<b>Post-translational modifications</b>	<p>Proteolytic processing produces the C-terminal angiogenic peptide, endorepellin. This peptide can be further processed to produce the LG3 peptide.</p> <p>N- and O-glycosylated; contains three heparan sulfate chains. The LG3 peptide contains at least three and up to five potential O-glycosylation sites but no N-glycosylation.</p>
<b>Cellular localization</b>	<p>Secreted &gt; extracellular space &gt; extracellular matrix &gt; basement membrane.</p>

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

---

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors