

Product datasheet

Human IFIH1 (MDA5) knockout A549 cell line ab266997

4 Images

Overview

Product name	Human IFIH1 (MDA5) knockout A549 cell line
Parental Cell Line	A549
Organism	Human
Mutation description	Knockout achieved by using CRISPR/Cas9, 1 bp deletion in exon 1 and 1 bp insertion in exon 1 and 8 bp deletion in exon 1
Passage number	<20
Knockout validation	Sanger Sequencing
Biosafety level	1
General notes	<p>Recommended control: Human wild-type A549 cell line (ab255450). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.</p> <p>Cryopreservation cell medium: Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.</p> <p>Culture medium: F-12K + 10% FBS</p> <p>Initial handling guidelines: Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.</p> <ol style="list-style-type: none"> 1. Thaw the vial in 37°C water bath approximately 1-2 minutes. 2. Transfer the cell suspension (0.8 ml) to a 15 ml/50 ml conical sterile polypropylene centrifuge tube containing 8.4 ml pre-warmed culture medium, wash vial with an additional 0.8 ml culture medium (total volume 10 ml) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 ml represents minimum recommended dilution. 20 ml represents maximum recommended dilution. 3. Resuspend the cell pellet in 5 ml pre-warmed culture medium and count using a haemocytometer (Click here to view haemocytometer protocol) or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of 2×10^4 cells/cm². This should allow for confluency within 48 hours. Seeding density is given as a guide only and should be scaled to align with individual lab schedules. 4. Incubate the culture at 37°C incubator with 5% CO₂. Cultures should be monitored daily. <p>Subculture guidelines:</p> <p>All seeding densities should be based on cell counts gained by established methods. A guide seeding density of 2×10^4 cells/cm² is recommended for confluency (80-90% confluence) within 48 hours.</p>

A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.

Cells should be passaged when they have achieved 80-90% confluence.

[Click here to view the Mammalian cell tissue culture protocol](#)

This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our [limited use license](#) and [patent pages](#).

Properties

Number of cells	1 x 10 ⁶ cells/vial, 1 mL
Viability	~90%
Adherent /Suspension	Adherent
Tissue	Lung
Cell type	epithelial
Disease	Carcinoma
Gender	Male
STR Analysis	Amelogenin X,YD5S818: 11 D13S317: 11 D7S820: 8, 11 D16S539: 11, 12 WWA: 14 TH01: 8,9.3 TPOX: 8,11 CSF1PO: 10, 12
Mycoplasma free	Yes
Storage instructions	Shipped on Dry Ice. Store in liquid nitrogen.
Storage buffer	Constituents: 8.7% DMSO, 2% Cellulose, methyl ether
Purity	Immunogen affinity purified

Target

Function	RNA helicase that, through its ATP-dependent unwinding of RNA, may function to promote message degradation by specific RNases. Seems to have growth suppressive properties. Involved in innate immune defense against viruses. Upon interaction with intracellular dsRNA produced during viral replication, triggers a transduction cascade involving MAVS/IPS1, which results in the activation of NF-kappa-B, IRF3 and IRF7 and the induction of the expression of antiviral cytokines such as IFN-beta and RANTES (CCL5). ATPase activity is specifically induced by dsRNA. Essential for the production of interferons in response to picornaviruses.
Tissue specificity	Widely expressed, at a low level. Expression is detected at slightly highest levels in placenta, pancreas and spleen and at barely levels in detectable brain, testis and lung.
Involvement in disease	Genetic variation in IFIH1 is associated with diabetes mellitus insulin-dependent type 19 (IDDM19) [MIM:610155]. A multifactorial disorder of glucose homeostasis that is characterized by susceptibility to ketoacidosis in the absence of insulin therapy. Clinical features are polydipsia, polyphagia and polyuria which result from hyperglycemia-induced osmotic diuresis and secondary thirst. These derangements result in long-term complications that affect the eyes, kidneys, nerves, and blood vessels. Note=IFIH1 is the CADM-140 autoantigen, involved in clinically amyopathic dermatomyositis (CADM). This is a chronic inflammatory disorder that shows typical skin manifestations of dermatomyositis but has no or little evidence of clinical myositis. Anti-CADM-140 antibodies appear to be specific to dermatomyositis, especially CADM. Patients with anti-CADM-140

antibodies frequently develop life-threatening acute progressive interstitial lung disease (ILD).

Sequence similarities

Belongs to the helicase family.
Contains 2 CARD domains.
Contains 1 helicase ATP-binding domain.
Contains 1 helicase C-terminal domain.

Post-translational modifications

During apoptosis, processed into 3 cleavage products. The helicase-containing fragment, once liberated from the CARD domains, translocate from the cytoplasm to the nucleus. The processed protein significantly sensitizes cells to DNA degradation.

Cellular localization

Cytoplasm. Nucleus. May be found in the nucleus, during apoptosis.

Images

```
Mut GCTT CAGGGCCAGGGT GAAAAAT GT ACAT CC ----- CCT GT GCT GGACT ACCT GACCT
WT  GCTT CAGGGCCAGGGT GAAAAAT GT ACAT CCAAGT GGAGCCT GT GCT GGACT ACCT GACCT
```

Sanger Sequencing - Human IFIH1 knockout A549 cell line (ab266997)

Allele-1: 8 bp deletion in exon1

```
Mut GCTT CAGGGCCAGGGT GAAAAAT GT ACAT CC - GGT GGAGCCT GT GCT GGACT ACCT GACCT
WT  GCTT CAGGGCCAGGGT GAAAAAT GT ACAT CCAAGT GGAGCCT GT GCT GGACT ACCT GACCT
```


Sanger Sequencing - Human IFIH1 knockout A549 cell line (ab266997)

Allele-2: 1 bp deletion in exon 1.

```
Mut GCTT CAGGGCCAGGGT GAAAAAT GT ACAT CCAAGT GGAGCCT GT GCT GGACT ACCT GACC
WT  GCTT CAGGGCCAGGGT GAAAAAT GT ACAT CCAAGT GGAGCCT GT GCT GGACT ACCT GACC
```

Sanger Sequencing - Human IFIH1 knockout A549 cell line (ab266997)

Allele-3: 1 bp insertion in exon 1.



Human IFIH1 (MDA5) knockout A549 cell line (ab266997)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors