

Product datasheet

Human IRF2 knockout A549 cell line ab267016

3 Images

Overview

Product name	Human IRF2 knockout A549 cell line
Parental Cell Line	A549
Organism	Human
Mutation description	Knockout achieved by using CRISPR/Cas9, 14 bp deletion in exon 4 and 1 bp insertion in exon 4
Passage number	<20
Knockout validation	Sanger Sequencing
Biosafety level	1
General notes	<p>Recommended control: Human wild-type A549 cell line (ab255450). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.</p> <p>Cryopreservation cell medium: Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.</p> <p>Culture medium: F-12K + 10% FBS</p> <p>Initial handling guidelines: Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.</p> <ol style="list-style-type: none"> 1. Thaw the vial in 37°C water bath approximately 1-2 minutes. 2. Transfer the cell suspension (0.8 ml) to a 15 ml/50 ml conical sterile polypropylene centrifuge tube containing 8.4 ml pre-warmed culture medium, wash vial with an additional 0.8 ml culture medium (total volume 10 ml) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 ml represents minimum recommended dilution. 20 ml represents maximum recommended dilution. 3. Resuspend the cell pellet in 5 ml pre-warmed culture medium and count using a haemocytometer (Click here to view haemocytometer protocol) or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of 2×10^4 cells/cm². This should allow for confluency within 48 hours. Seeding density is given as a guide only and should be scaled to align with individual lab schedules. 4. Incubate the culture at 37°C incubator with 5% CO₂. Cultures should be monitored daily. <p>Subculture guidelines:</p> <p>All seeding densities should be based on cell counts gained by established methods. A guide seeding density of 2×10^4 cells/cm² is recommended for confluency (80-90% confluence) within 48 hours.</p> <p>A partial media change 24 hours prior to subculture may be helpful to encourage growth, if</p>

required.

Cells should be passaged when they have achieved 80-90% confluence.

[Click here to view the Mammalian cell tissue culture protocol](#)

This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our [limited use license](#) and [patent pages](#).

Properties

Number of cells	1 x 10 ⁶ cells/vial, 1 mL
Viability	~90%
Adherent /Suspension	Adherent
Tissue	Lung
Cell type	epithelial
Disease	Carcinoma
Gender	Male
STR Analysis	Amelogenin X,YD5S818: 11 D13S317: 11 D7S820: 8, 11 D16S539: 11, 12 WWA: 14 TH01: 8,9,3 TPOX: 8,11 CSF1PO: 10, 12
Mycoplasma free	Yes
Storage instructions	Shipped on Dry Ice. Store in liquid nitrogen.
Storage buffer	Constituents: 8.7% DMSO, 2% Cellulose, methyl ether
Purity	Immunogen affinity purified

Target

Function	Specifically binds to the upstream regulatory region of type I IFN and IFN-inducible MHC class I genes (the interferon consensus sequence (ICS)) and represses those genes. Also acts as an activator for several genes including H4 and IL7. Constitutively binds to the ISRE promoter to activate IL7. Involved in cell cycle regulation through binding the site II (HiNF-M) promoter region of H4 and activating transcription during cell growth. Antagonizes IRF1 transcriptional activation.
Tissue specificity	Expressed throughout the epithelium of the colon. Also expressed in lamina propria.
Sequence similarities	Belongs to the IRF family. Contains 1 IRF tryptophan pentad repeat DNA-binding domain.
Post-translational modifications	Acetylated by CBP/ p300 during cell-growth. Acetylation on Lys-75 is required for stimulation of H4 promoter activity. The major sites of sumoylation are Lys-137 and Lys-293. Sumoylation by SUMO1 increases its transcriptional repressor activity on IRF1 and diminishes its ability to activate ISRE and H4 promoter.
Cellular localization	Nucleus.

Images

Mut	TT CAGGGT CT ACC GAAT GCT GCCCT ----- CT AAGAAAGGT AAAGGCAT T
WT	TT CAGGGT CT ACC GAAT GCT GCCCT AT CAGAACGGCCTT CT AAGAAAGGT AAAGGCAT T


Allele-1: 14 bp deletion in exon4

Sanger Sequencing - Human IRF2 knockout A549
cell line (ab267016)

Mut	TT CAGGGT CT ACC GAAT GCT GCCCT CAT CAGAACGGCCTT CT AAGAAAGGT AAAGGCAT
WT	TT CAGGGT CT ACC GAAT GCT GCCCT AT CAGAACGGCCTT CT AAGAAAGGT AAAGGCAT

Allele-2: 1 bp insertion in exon 4.

Sanger Sequencing - Human IRF2 knockout A549
cell line (ab267016)



Human IRF2 knockout A549 cell line (ab267016)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors