

## Product datasheet

# Anti-Integrin alpha 5 antibody [5H10-27 (MFR5)] ab25251

★★★★★ 2 Abreviews 8 References

### Overview

---

<b>Product name</b>	Anti-Integrin alpha 5 antibody [5H10-27 (MFR5)]
<b>Description</b>	Rat monoclonal [5H10-27 (MFR5)] to Integrin alpha 5
<b>Host species</b>	Rat
<b>Specificity</b>	ab25251 is specific for the alpha 5 subunit of the integrin alpha 5 beta 1 fibronectin receptor.
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse
<b>Immunogen</b>	Tissue, cells or virus corresponding to Mouse Integrin alpha 5. C57BL/6 x A/J F1 mouse mast cell line MC/9

### General notes

Soluble ab25251 inhibits VLA5-mediated functions in vitro. Immobilized ab25251 has also been demonstrated to co-stimulate the proliferative response of CD8+ T cells to plate-bound antiCD3 monoclonal antibody.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

### Properties

---

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
<b>Storage buffer</b>	pH: 8.20 Constituent: 100% Borate buffered saline
<b>Primary antibody notes</b>	Soluble ab25251 inhibits VLA5-mediated functions in vitro. Immobilized ab25251 has also been demonstrated to co-stimulate the proliferative response of CD8+ T cells to plate-bound antiCD3 monoclonal antibody.

<b>Clonality</b>	Monoclonal
<b>Clone number</b>	5H10-27 (MFR5)
<b>Isotype</b>	IgG2a
<b>Light chain type</b>	kappa

---

## Target

<b>Function</b>	Integrin alpha-5/beta-1 is a receptor for fibronectin and fibrinogen. It recognizes the sequence R-G-D in its ligands. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions.
<b>Sequence similarities</b>	Belongs to the integrin alpha chain family. Contains 7 FG-GAP repeats.
<b>Cellular localization</b>	Membrane.

---

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

---

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

---

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors