

Product datasheet

MCF7 whole cell lysate ab29537

Overview

Product name MCF7 whole cell lysate

General notes

Protein is isolated from whole tissue homogenates using a proprietary technique. The tissue was frozen in liquid nitrogen immediately after excision and then stored at -70°C. Quality control: 1. The isolated protein pattern on SDS-PAGE gel is visualized by coomassie blue staining. The pattern is consistent with each lot. 2. The isolated protein is Western Blot analyzed by either a GAPDH or beta-actin antibody. The expression level is consistent with each lot.

Abcam has not and does not intend to apply for the REACH Authorisation of customers' uses of products that contain European Authorisation list (Annex XIV) substances.

It is the responsibility of our customers to check the necessity of application of REACH Authorisation, and any other relevant authorisations, for their intended uses.

Tested applications **Suitable for:** WB

Properties

Mycoplasma free Yes

Form Liquid

Storage instructions Store at -80°C. Allow to warm to room temp and agitate gently before aliquotting.

Storage buffer Constituents: 1% 4-Nonylphenol, branched, ethoxylated, Potassium chloride, 0.1% Bromophenol blue, Magnesium chloride, 1.2% Beta mercaptoethanol, 2% Sodium lauryl sulfate, HEPES, Sucrose, 10% Glycerol

With Protease inhibitor cocktail/EDTA/Sodium deoxycholate

Lysate notes Protein is isolated from whole tissue homogenates using a proprietary technique. The tissue was frozen in liquid nitrogen immediately after excision and then stored at -70°C. Quality control: 1. The isolated protein pattern on SDS-PAGE gel is visualized by coomassie blue staining. The pattern is consistent with each lot. 2. The isolated protein is Western Blot analyzed by either a GAPDH or beta-actin antibody. The expression level is consistent with each lot.

Background MCF-7 cells are a well-characterized estrogen receptor (ER) positive control cell line (cells are positive for cytoplasmic estrogen receptors) and therefore are a useful in vitro model of breast cancer to study the role of estrogen in breast cancer. Cells are also positive for cytokeratin and negative for desmin , endothelin , GFAP , neurofilament, vimentin.

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab29537 in the following tested applications. The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		Use at an assay dependent dilution. Samples need to be boiled for 5 minutes and put on ice before use.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors