

Product datasheet

Anti-Menin antibody [EPR3986] ab92443

KO VALIDATED Recombinant RabMAb

5 References 9 Images

Overview

Product name	Anti-Menin antibody [EPR3986]
Description	Rabbit monoclonal [EPR3986] to Menin
Host species	Rabbit
Tested applications	Suitable for: WB, IP, IHC-P, ICC Unsuitable for: Flow Cyt
Species reactivity	Reacts with: Mouse, Rat, Human
Immunogen	Synthetic peptide within Human Menin aa 600-700 (C terminal). The exact sequence is proprietary.
Positive control	WB: Wild-type HAP1, Jurkat cell lysates. IP: Jurkat whole and MEF cell lysates. ICC: Jurkat cells
General notes	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 0.05% BSA, 40% Glycerol (glycerin, glycerine), 59% PBS

Purity	Protein A purified
Clonality	Monoclonal
Clone number	EPR3986
Isotype	IgG

Applications

The Abpromise guarantee Our [Abpromise guarantee](#) covers the use of ab92443 in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB		1/1000. Predicted molecular weight: 68 kDa. For unpurified use between 1/10,000-1/50,000.
IP		1/10 - 1/100.
IHC-P		1/100 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. For unpurified use between 1/100-1/250.
ICC		1/50. For unpurified use between 1/250-1/500.

Application notes Is unsuitable for Flow Cyt.

Target

Function Essential component of a MLL/SET1 histone methyltransferase (HMT) complex, a complex that specifically methylates 'Lys-4' of histone H3 (H3K4). Functions as a transcriptional regulator. Binds to the TERT promoter and represses telomerase expression. Plays a role in TGFB1-mediated inhibition of cell-proliferation, possibly regulating SMAD3 transcriptional activity. Represses JUND-mediated transcriptional activation on AP1 sites, as well as that mediated by NFKB subunit RELA. Positively regulates HOXC8 and HOXC6 gene expression. May be involved in normal hematopoiesis through the activation of HOXA9 expression (By similarity). May be involved in DNA repair.

Tissue specificity Ubiquitous.

Involvement in disease Defects in MEN1 are the cause of familial multiple endocrine neoplasia type I (MEN1) [MIM:131100]. Autosomal dominant disorder characterized by tumors of the parathyroid glands, gastro-intestinal endocrine tissue, the anterior pituitary and other tissues. Cutaneous lesions and nervous-tissue tumors can exist. Prognosis in MEN1 patients is related to hormonal hypersecretion by tumors, such as hypergastrinemia causing severe peptic ulcer disease (Zollinger-Ellison syndrome, ZES), primary hyperparathyroidism, and acute forms of hyperinsulinemia.

Defects in MEN1 are the cause of familial isolated hyperparathyroidism (FIHP) [MIM:145000]; also known as hyperparathyroidism type 1 (HRPT1). FIHP is an autosomal dominant disorder characterized by hypercalcemia, elevated parathyroid hormone (PTH) levels, and uniglandular or

multiglandular parathyroid tumors.


Phosphorylated upon DNA damage, probably by ATM or ATR.

Post-translational modifications

Cellular localization


Nucleus. Concentrated in nuclear body-like structures. Relocates to the nuclear matrix upon gamma irradiation.

Images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat cerebrum tissue sections labelling Menin with purified ab92443 at 1/100 dilution (6.08 µg/ml). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1/0 dilution. PBS instead of the primary antibody was used as the negative control.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Menin antibody [EPR3986] (ab92443)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue sections labelling Menin with purified ab92443 at 1/100 dilution (6.08 µg/ml). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1/0 dilution. PBS instead of the primary antibody was used as the negative control.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Menin antibody [EPR3986] (ab92443)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Menin antibody [EPR3986] (ab92443)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human testis tissue sections labelling Menin with purified ab92443 at 1/500 dilution (1.22 µg/ml). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1/0 dilution. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-Menin antibody [EPR3986] (ab92443)

Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: Menin knockout HAP1 cell lysate (20 µg)

Lane 3: Jurkat cell lysate (20 µg)

Lane 4: A431 cell lysate (20 µg)


Lanes 1 - 4: Merged signal (red and green). Green - ab92443 observed at 74 kDa. Red - loading control, ab8245, observed at 37 kDa.

ab92443 was shown to recognize Menin when Menin knockout samples were used, along with additional cross-reactive bands. Wild-type and Menin knockout samples were subjected to SDS-PAGE. ab92443 and ab8245 (loading control to GAPDH) were diluted 1/10,000 and 1/2000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.



Immunocytochemistry - Anti-Menin antibody
[EPR3986] (ab92443)

Immunocytochemistry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling Menin with purified ab92443 at 1/50 dilution (12.1 µg/mL). Cells were fixed in 100% Methanol and permeabilized with 0.1% tritonX-100. Cells were counterstained with [ab195889](#) Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.6 µg/mL). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150078](#)) was used as the secondary antibody at 1/1000 (2 µg/mL) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunoprecipitation - Anti-Menin antibody
[EPR3986] (ab92443)

Purified ab92443 at 1/50 dilution (2µg) immunoprecipitating Menin in Jurkat whole cell lysate.

Lane 1 (input): Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate 10µg

Lane 2 (+): ab92443 + Jurkat whole cell lysate.


Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab92443 in Jurkat whole cell lysate.

VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) (1/1000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm/TBST.

Observed band size: 75 kDa



Immunoprecipitation - Anti-Menin antibody
[EPR3986] (ab92443)

Menin was immunoprecipitated from 0.35 mg MEF (mouse embryonic fibroblast (immortalized)) whole cell lysate 10 ug with ab92443 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab92443 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.


Lane 1: MEF (mouse embryonic fibroblast (immortalized)) whole cell lysate 10 µg

Lane 2: ab92443 IP in MEF whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab92443 in MEF whole cell lysate

Blocking and dilution buffer and concentration: 5% NDM/TBST.

Exposure time: 3 minutes



Western blot - Anti-Menin antibody [EPR3986]
(ab92443)

All lanes : Anti-Menin antibody [EPR3986] (ab92443) at 1/1000 dilution (Purified)

Lane 1 : Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate

Lane 2 : MEF (Mouse embryonic fibroblast (immortalized)) whole cell lysate

Lane 3 : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysate

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 68 kDa

Observed band size: 75 kDa

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Menin antibody [EPR3986] (ab92443)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors