

Product datasheet

Anti-MYO6 antibody ab106288

Overview

Product name	Anti-MYO6 antibody
Description	Rabbit polyclonal to MYO6
Host species	Rabbit
Tested applications	Suitable for: IHC-P, IHC-Fr, WB, IP
Species reactivity	Reacts with: Human Predicted to work with: a wide range of other species ▲
Immunogen	His tagged Human MYO6.

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Storage buffer	Constituent: Whole serum
Purity	Whole antiserum
Clonality	Polyclonal
Isotype	IgG

Applications

Our [Abpromise guarantee](#) covers the use of **ab106288** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
IHC-P		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.
WB		1/500 - 1/1000. Predicted molecular weight: 150 kDa.
IP		1/2.

Target

Function	Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Myosin 6 is a reverse-direction motor protein that moves towards the minus-end of actin filaments. Has slow rate of actin-activated ADP release due to weak ATP binding. Functions in a variety of intracellular processes such as vesicular membrane trafficking and cell migration. Required for the structural integrity of the Golgi apparatus via the p53-dependent pro-survival pathway. Appears to be involved in a very early step of clathrin-mediated endocytosis in polarized epithelial cells. May act as a regulator of F-actin dynamics. May play a role in transporting DAB2 from the plasma membrane to specific cellular targets. Required for structural integrity of inner ear hair cells.
Tissue specificity	Expressed in most tissues examined including heart, brain, placenta, pancreas, spleen, thymus, prostate, testis, ovary, small intestine and colon. Highest levels in brain, pancreas, testis and small intestine. Also expressed in fetal brain and cochlea. Isoform 1 and isoform 2, containing the small insert, and isoform 4, containing neither insert, are expressed in unpolarized epithelial cells.
Involvement in disease	Defects in MYO6 are the cause of deafness autosomal dominant type 22 (DFNA22) [MIM:606346]. DFNA22 is a form of sensorineural hearing loss. Sensorineural deafness results from damage to the neural receptors of the inner ear, the nerve pathways to the brain, or the area of the brain that receives sound information. DFNA22 is progressive and postlingual, with onset during childhood. By the age of approximately 50 years, affected individuals invariably have profound sensorineural deafness. Defects in MYO6 are the cause of deafness autosomal recessive type 37 (DFNB37) [MIM:607821]. Defects in MYO6 are the cause of deafness sensorineural with hypertrophic cardiomyopathy (DFNHCM) [MIM:606346].
Sequence similarities	Contains 1 IQ domain. Contains 1 myosin head-like domain.
Domain	Divided into three regions: a N-terminal motor (head) domain, followed by a neck domain consisting of a calmodulin-binding linker domain and a single IQ motif, and a C-terminal tail region with a coiled-coil and a unique globular domain required for interaction with other proteins.
Post-translational modifications	Phosphorylation in the motor domain, induced by EGF, results in translocation of MYO6 from the cell surface to membrane ruffles and affects F-actin dynamics. Phosphorylated in vitro by p21-activated kinase (PAK).
Cellular localization	Cytoplasmic vesicle > clathrin-coated vesicle membrane; Cytoplasmic vesicle > clathrin-coated vesicle membrane. Cell projection > ruffle membrane and Golgi apparatus > trans-Golgi network membrane. Golgi apparatus. Nucleus. Cytoplasm > perinuclear region. Membrane > clathrin-coated pit. Cell projection > ruffle membrane. Also present in endocytic vesicles, and membrane ruffles. Translocates from membrane ruffles, endocytic vesicles and cytoplasm to Golgi apparatus, perinuclear membrane and nucleus through induction by p53 and p53-induced DNA damage. Recruited into membrane ruffles from cell surface by EGF-stimulation. Colocalizes with DAB2 in clathrin-coated pits/vesicles. Colocalizes with OPTN at the Golgi complex and in vesicular structures close to the plasma membrane.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors